

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau




(43) International Publication Date
27 May 2004 (27.05.2004)

PCT

(10) International Publication Number
WO 2004/043953 A1

- (51) International Patent Classification⁷: **C07D 417/04**, A61K 31/506, A61P 35/00
- (74) Agents: **CLYDE-WATSON, Zoe et al.**; D Young & Co., 21 New Fetter Lane, London EC4A 1DA (GB).
- (21) International Application Number:
PCT/GB2003/004973
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (22) International Filing Date:
14 November 2003 (14.11.2003)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
0226583.3 14 November 2002 (14.11.2002) GB
- (84) Designated States (*regional*): ARIPO patent (BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- (71) Applicant (*for all designated States except US*): **CYCLACEL LIMITED** [GB/GB]; 12 St. James's Square, London SW1Y 4RB (GB).
- (72) Inventors; and
- (75) Inventors/Applicants (*for US only*): **WANG, Shudong** [AU/GB]; Burnside Mill, Forfar, Angus DD8 2RZ (GB). **MEADES, Christopher** [GB/GB]; 32 Chirnside Place, Whitehazel Park, Dundee DD4 0TE (GB). **WOOD, Gavin** [GB/GB]; Whinrig, Millbank, Cupar, Fife KY15 5DP (GB). **O'BOYLE, Janice** [GB/GB]; 11 Gullane Terrace, Dundee DD2 3BT (GB). **MCINNES, Campbell** [GB/GB]; 8 Mary Findlay Drive, Longforgan, Dundee DD2 5JE (GB). **FISCHER, Peter** [CH/GB]; Denley Lodge, 1 Arbirlot Road, Arbroath, Angus DD11 2EN (GB).
- Declaration under Rule 4.17:**
— *of inventorship (Rule 4.17(iv)) for US only*
- Published:**
— *with international search report*
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: PYRIMIDINE COMPOUNDS



(I)

(57) Abstract: The present invention relates to substituted pyrimidines of formula (I), their preparation, pharmaceutical compositions containing them and their use as inhibitors of cyclin-dependent kinases (CDKs) and hence their use in the treatment of proliferative disorders and/or viral disorders.

WO 2004/043953 A1

PYRIMIDINE COMPOUNDS

The present invention relates to new 2-substituted-4-heteroaryl-pyrimidine derivatives and their use in therapy. More specifically, the invention relates to 2-substituted-4-heteroaryl-pyrimidine derivatives having improved solubility properties.


BACKGROUND

We have previously disclosed 2-substituted-4-heteroaryl-pyrimidines and their use in the treatment of proliferative disorders (Fischer PM, Wang S. PCT Intl. Patent Appl. Publ. WO 01/072745; Cyclacel Limited, UK, 2001). These compounds inhibit cyclin-dependent protein kinases (CDKs), in particular CDK4 / cyclin D, CDK2 / cyclin E, CDK2 / cyclin A, and CDK1 / cyclin B, *i.e.* enzyme complexes that are important in human cell cycle progression. Furthermore, 2-phenylamino-4-heteroaryl-pyrimidines possess selective *in vitro* and *in vivo* antiproliferative activity against a range of human tumour cells (Wang S, Blake D, Clarke R, Duff S, McClue SJ, McInnes C, Melville J, Stewart K, Taylor P, Westwood R, Wood G, Wu S-Y, Zhelev NZ, Zheleva DI, Walkinshaw M, Lane DP, Fischer PM. Proc. Amer. Assoc. Cancer Res. 2002; 43: 4202).

The present invention seeks to provide further 2-substituted-4-heteroaryl-pyrimidines. More specifically, the present invention preferably seeks to provide 2-substituted-4-heteroaryl-pyrimidines which display improved aqueous solubility and/or bioavailability.

STATEMENT OF INVENTION

A first aspect of the invention relates to a compound of formula I, or a pharmaceutically acceptable salt thereof,



I

wherein:

- (A) one of X^1 and X^2 is S, and the other of X^1 and X^2 is N;
 5 "a" is a single bond;
 "b", "c", "d", "e" and "f" are single or double bonds so as to form a thiazolyl ring;
 R^2 is independently as defined below for R^1 , R^{3-8} ; or
 (B) one of X^1 and X^2 is S, and the other of X^1 and X^2 is NR^{17} ;
 "a" and "d" are each double bonds;
 10 "b", "c", "e" and "f" are each single bonds;
 R^2 is oxo; and
 R^{17} is H or alkyl;

where:

- 15 Z is NH, NHCO, NHSO₂, NHCH₂, CH₂, CH₂CH₂, CH=CH, SO₂, or SO;

- R^1 , R^3 , R^4 , R^5 , R^6 , R^7 and R^8 are each independently H, alkyl, alkyl- R^9 , aryl, aryl- R^9 , aralkyl, aralkyl- R^9 , halogeno, NO₂, CN, OH, O-alkyl, COR⁹, COOR⁹, O-aryl, O- R^9 , NH₂, NH-alkyl, NH-aryl, N-(alkyl)₂, N-(aryl)₂, N-(alkyl)(aryl), NH- R^9 , N-(R^9)(R^{10}), N-(alkyl)(R^9), N-(aryl)(R^9), COOH, CONH₂, CONH-alkyl, CONH-aryl, CON-(alkyl)(R^9), CON(aryl)(R^9), CONH- R^9 , CON-(R^9)(R^{10}), SO₃H, SO₂-alkyl, SO₂-alkyl- R^9 , SO₂-aryl, SO₂-aryl- R^9 , SO₂NH₂, SO₂NH- R^9 , SO₂N-(R^9)(R^{10}), CF₃, CO-alkyl, CO-alkyl- R^9 , CO-aryl, CO-aryl- R^9 or R^{11} , wherein alkyl, aryl, aralkyl groups may be further substituted with one

3

or more groups selected from halogeno, NO₂, OH, O-methyl, NH₂, COOH, CONH₂ and CF₃;

5 wherein at least one of R¹, R², R³, R⁴, R⁵, R⁶, R⁷ and R⁸ is an R⁹ or R¹⁰-containing group, or is R¹¹;

R⁹ and R¹⁰ are each independently solubilising groups selected from:

- (i) - a mono-, di- or polyhydroxylated alicyclic group;
- a di- or polyhydroxylated aliphatic or aromatic group;
10 - a carbohydrate derivative;
- an O- and/or S-containing heterocyclic group optionally substituted by one or more hydroxyl groups;
- an aliphatic or aromatic group containing a carboxamide, sulfoxide, sulfone, or sulfonamide function; or
15 - a halogenated alkylcarbonyl group;
- (ii) COOH, SO₃H, OSO₃H, PO₃H₂, or OPO₃H₂;
- (iii) Y, where Y is selected from an alicyclic, aromatic, or heterocyclic group comprising one or more of the functions =N-, -O-, -NH₂, -NH-, a quarternary amine salt, guanidine, and amidine, where Y is optionally substituted by one or more
20 substituents selected from:
- SO₂-alkyl;
 - alkyl optionally substituted by one or more OH groups;
 - CO-alkyl;
 - aralkyl;
 - 25 - COO-alkyl; and
 - an ether group optionally substituted by one or more OH groups; and
- where Y is other than pyridinyl;
- (iv) a natural or unnatural amino acid, a peptide or a peptide derivative;

each R^{11} is a solubilising group as defined for R^9 and R^{10} in (i) or (iv) above; or is selected from:

- (v) OSO_3H , PO_3H_2 , or OPO_3H_2 ;
- (vi) Y as defined above, but excluding guanidine and quarternary amine salts;
- 5 (vii) $NHCO(CH_2)_m[NHCO(CH_2)_{m'}]_p[NHCO(CH_2)_{m''}]_qY$ or $NHCO(CH_2)_tNH(CH_2)_{t'}Y$ where p and q are each 0 or 1, and m, m', m'', t and t' are each independently an integer from 1 to 10; and
- (viii) $(CH_2)_nNR^{14}COR^{12}$, $(CH_2)_nNR^{15}SO_2R^{13}$, or SO_2R^{16} , where R^{12} , R^{13} and R^{16} are each alkyl groups optionally comprising one or more heteroatoms, and which are optionally substituted by one or more substituents selected from OH, NH_2 , halogen and NO_2 , R^{14} and R^{15} are each independently H or alkyl, and n and n' are each independently 0, 1, 2, or 3;
- (ix) an ether or polyether optionally substituted by one or more hydroxyl groups or one or more Y groups;
- 15 (x) $(CH_2)_rNH_2$; where r is 0, 1, 2, or 3;
- (xi) $(CH_2)_{r'}OH$; where r' is 0, 1, 2, or 3;

with the proviso that the compound is other than:

- 2-Chloro-N-{4-methyl-5-[2-(3-nitro-phenylamino)-pyrimidin-4-yl]-thiazol-2-yl}-acetamide;
- 20 N-{4-Methyl-5-[2-(3-nitro-phenylamino)-pyrimidin-4-yl]-thiazol-2-yl}-methane sulfonamide;
- 2-Chloro-N-{5-[2-(4-fluoro-phenylamino)-pyrimidin-4-yl]-4-methyl-thiazol-2-yl}-acetamide;
- 25 {3-[4-(2,4-Dimethyl-thiazol-5-yl)-pyrimidin-2-ylamino]-phenyl}-methanol;
- 2-{4-[4-(2-Amino-4-methyl-thiazol-5-yl)-pyrimidin-2-ylamino]-phenyl}-ethanol; or
- 2-{4-[4-(2,4-Dimethyl-thiazol-5-yl)-pyrimidin-2-ylamino]-phenyl}-ethanol.


5

A second aspect of the invention relates to a pharmaceutical composition comprising a compound of formula I as defined above admixed with a pharmaceutically acceptable diluent, excipient or carrier.

- 5 A third aspect of the invention relates to the use of a compound of formula I as defined above in the preparation of a medicament for treating a proliferative disorder.

A fourth aspect of the invention relates to the use of a compound of formula Ia, or a pharmaceutically acceptable salt thereof,

10



Ia

wherein:

- (A) one of X¹ and X² is S, and the other of X¹ and X² is N;
 15 "a" is a single bond;
 "b", "c", "d", "e" and "f" are single or double bonds so as to form a thiazolyl ring;
 R² is independently as defined below for R¹, R³⁻⁸; or
 (B) one of X¹ and X² is S, and the other of X¹ and X² is NR¹⁷;
 "a" and "d" are each double bonds; and
 20 "b", "c", "e" and "f" are each single bonds;
 R² is oxo;
 R¹⁷ is H or alkyl;

where:

6

Z is NH, NHCO, NHSO₂, NHCH₂, CH₂, CH₂CH₂, CH=CH, SO₂, or SO;

R¹, R³, R⁴, R⁵, R⁶, R⁷ and R⁸ are each independently H, alkyl, alkyl-R⁹, aryl, aryl-R⁹, aralkyl, aralkyl-R⁹, halogeno, NO₂, CN, OH, O-alkyl, COR⁹, COOR⁹, O-aryl, O-R⁹, NH₂,
 5 NH-alkyl, NH-aryl, N-(alkyl)₂, N-(aryl)₂, N-(alkyl)(aryl), NH-R⁹, N-(R⁹)(R¹⁰), N-(alkyl)(R⁹), N-(aryl)(R⁹), COOH, CONH₂, CONH-alkyl, CONH-aryl, CON-(alkyl)(R⁹), CON(aryl)(R⁹), CONH-R⁹, CON-(R⁹)(R¹⁰), SO₃H, SO₂-alkyl, SO₂-alkyl-R⁹, SO₂-aryl, SO₂-aryl-R⁹, SO₂NH₂, SO₂NH-R⁹, SO₂N-(R⁹)(R¹⁰), CF₃, CO-alkyl, CO-alkyl-R⁹, CO-aryl, CO-aryl-R⁹ or R¹¹, wherein alkyl, aryl, aralkyl groups may be further substituted with one
 10 or more groups selected from halogeno, NO₂, OH, O-methyl, NH₂, COOH, CONH₂ and CF₃;

wherein at least one of R¹, R², R³, R⁴, R⁵, R⁶, R⁷ and R⁸ is an R⁹ or R¹⁰-containing group, or is R¹¹;

15

R⁹ and R¹⁰ are each independently solubilising groups selected from:

- (i) - a mono-, di- or polyhydroxylated alicyclic group;
 - a di- or polyhydroxylated aliphatic or aromatic group;
 - a carbohydrate derivative;
 20 - an O- and/or S-containing heterocyclic group optionally substituted by one or more hydroxyl groups;
 - an aliphatic or aromatic group containing a carboxamide, sulfoxide, sulfone, or sulfonamide function; or
 - a halogenated alkylcarbonyl group;
- 25 (ii) COOH, SO₃H, OSO₃H, PO₃H₂, or OPO₃H₂;
- (iii) Y, where Y is selected from an alicyclic, aromatic, or heterocyclic group comprising one or more of the functions =N-, -O-, -NH₂, -NH-, a quarternary amine salt, guanidine, and amidine, where Y is optionally substituted by one or more substituents selected from:
 30 - SO₂-alkyl;

7

- alkyl optionally substituted by one or more OH groups;
- CO-alkyl;
- aralkyl;
- COO-alkyl; and

5 - an ether group optionally substituted by one or more OH groups; and

where Y is other than pyridinyl;

(iv) a natural or unnatural amino acid, a peptide or a peptide derivative;

10 each R^{11} is a solubilising group as defined for R^9 and R^{10} in (i) or (iv) above; or is selected from:

(v) OSO_3H , PO_3H_2 , or OPO_3H_2 ;

(vi) Y as defined above, but excluding guanidine and quaternary amine salts;

(vii) $NHCO(CH_2)_m[NHCO(CH_2)_{m'}]_p[NHCO(CH_2)_{m''}]_qY$ or $NHCO(CH_2)_tNH(CH_2)_{t'}Y$
 15 where p and q are each 0 or 1, and m, m', m'', t and t' are each independently an integer from 1 to 10; and

(viii) $(CH_2)_nNR^{14}COR^{12}$, $(CH_2)_nNR^{15}SO_2R^{13}$, or SO_2R^{16} , where R^{12} , R^{13} and R^{16} are
 20 each alkyl groups optionally comprising one or more heteroatoms, and which are optionally substituted by one or more substituents selected from OH, NH_2 , halogen and NO_2 , R^{14} and R^{15} are each independently H or alkyl, and n and n' are each independently 0, 1, 2, or 3;

(ix) an ether or polyether optionally substituted by one or more hydroxyl groups or one or more Y groups;

(x) $(CH_2)_rNH_2$; where r is 0, 1, 2, or 3;

(xi) $(CH_2)_{r'}OH$; where r' is 0, 1, 2, or 3;

25 in the preparation of a medicament for treating a viral disorder.

A fifth aspect of the invention relates to the use of a compound of formula I as defined above for inhibiting a protein kinase.

A sixth aspect of the invention relates to the use of a compound of formula I as defined above in an assay for identifying further candidate compounds capable of inhibiting a cyclin dependent kinase.

5 DETAILED DESCRIPTION

As used herein the term "alkyl" includes both straight chain and branched alkyl groups having from 1 to 8 carbon atoms, e.g. methyl, ethyl propyl, isopropyl, butyl, isobutyl, tert-butyl, pentyl, hexyl etc. and the term "lower alkyl" is similarly used for groups having from 1 to 4 carbon atoms.

10

As used herein, the term "aryl" refers to a substituted (mono- or poly-) or unsubstituted monoaromatic or polyaromatic system, wherein said polyaromatic system may be fused or unfused. Preferably, the term "aryl" includes groups having from 6 to 10 carbon atoms, e.g. phenyl, naphthyl etc. The term "aryl" is synonymous with the term "aromatic".

15

The term "aralkyl" is used as a conjunction of the terms alkyl and aryl as given above.

The term "alicyclic" refers to a cyclic aliphatic group.

20 The term "aliphatic" takes its normal meaning in the art and includes non-aromatic groups such as alkanes, alkenes and alkynes and substituted derivatives thereof.

As used herein, the term "carbohydrate derivative" refers to a compound of general formula $C_x(H_2O)_y$ or a derivative thereof. Preferably, the carbohydrate is a mono-, di- or tri-saccharide. Monosaccharides can exist as either straight chain or ring-shaped molecules and are classified according to the number of carbon atoms they possess; *trioses* have three carbons, *tetroses* four, *pentoses* five and *hexoses* six. Each of these subgroups may be further divided into aldoses and ketoses, depending on whether the molecule contains an aldehyde group (-CHO) or a ketone group (C=O). Typical examples of monosaccharides include glucose, fructose, and galactose. Disaccharides consist of two linked

30

9

monosaccharide molecules, and include for example, maltose and lactose. Trisaccharides consist of three linked monosaccharide molecules.

The term "derivative" as used herein includes chemical modification of an entity.


- 5 Illustrative of such chemical modifications would be replacement of hydrogen by a halo group, an alkyl group, an acyl group or an amino group.

The term "heterocycle" refers to a saturated or unsaturated cyclic group containing one or more heteroatoms in the ring.

10

As used herein the phrase "preparation of a medicament" includes the use of a compound of formula I directly as the medicament in addition to its use in a screening programme for further anti-viral agents or in any stage of the manufacture of such a medicament.


- 15 In one preferred embodiment, the invention relates to compounds of formula Ib, or pharmaceutically acceptable salts thereof,



- 20 wherein one of X^1 and X^2 is S, and the other of X^1 and X^2 is N, and Z, R^{1-8} are as defined above.

In another preferred embodiment, the invention relates to compounds of formula Ic, or pharmaceutically acceptable salts thereof,

25



wherein

one of X^1 and X^2 is S, and the other of X^1 and X^2 is N;

5

Z is NH, NHCO, NHSO₂, NHCH₂, CH₂, CH₂CH₂, CH=CH, SO₂, or SO;

10 R^1 , R^2 , R^3 , R^4 , R^5 , R^6 , R^7 and R^8 are each independently H, alkyl, alkyl- R^9 , aryl, aryl- R^9 , aralkyl, aralkyl- R^9 , halogeno, NO₂, CN, OH, O-alkyl, COR⁹, COOR⁹, O-aryl, O- R^9 , NH₂, NH-alkyl, NH-aryl, N-(alkyl)₂, N-(aryl)₂, N-(alkyl)(aryl), NH- R^9 , N-(R^9)(R^{10}), N-(alkyl)(R^9), N-(aryl)(R^9), COOH, CONH₂, CONH-alkyl, CONH-aryl, CON-(alkyl)(R^9), CON(aryl)(R^9), CONH- R^9 , CON-(R^9)(R^{10}), SO₃H, SO₂-alkyl, SO₂-alkyl- R^9 , SO₂-aryl, SO₂-aryl- R^9 , SO₂NH₂, SO₂NH- R^9 , SO₂N-(R^9)(R^{10}), CF₃, CO-alkyl, CO-alkyl- R^9 , CO-aryl, CO-aryl- R^9 or R^{11} , wherein alkyl, aryl, aralkyl groups may be further substituted with one
15 or more groups selected from halogeno, NO₂, OH, O-methyl, NH₂, COOH, CONH₂ and CF₃;

wherein at least one of R^1 , R^2 , R^3 , R^4 , R^5 , R^6 , R^7 and R^8 is an R^9 or R^{10} -containing group, or is R^{11} ;

20

R^9 and R^{10} are each independently solubilising groups selected from:

- (i) - a mono-, di- or polyhydroxylated alicyclic group;
- a di- or polyhydroxylated aliphatic or aromatic group;
- a carbohydrate derivative;

- an O- and/or S-containing heterocyclic group optionally substituted by one or more hydroxyl groups;
- an aliphatic or aromatic group containing a carboxamide, sulfoxide, sulfone, or sulfonamide function; or
- 5 - a halogenated alkylcarbonyl group;
- (ii) COOH , SO_3H , OSO_3H , PO_3H_2 , or OPO_3H_2 ;
- (iii) Y, where Y is selected from an alicyclic, aromatic, or heterocyclic group comprising one or more of the functions $=\text{N}-$, $-\text{O}-$, $-\text{NH}_2$, $-\text{NH}-$, a quarternary amine salt, guanidine, and amidine, where Y is optionally substituted by one or more
- 10 substituents selected from:
 - SO_2 -alkyl;
 - alkyl optionally substituted by one or more OH groups;
 - CO-alkyl;
 - aralkyl;
 - 15 - COO-alkyl; and
 - an ether group optionally substituted by one or more OH groups; and
 where Y is other than pyridinyl;
- (iv) a natural or unnatural amino acid, a peptide or a peptide derivative;
- 20 R^{11} is a solubilising group as defined for R^9 and R^{10} in (i) or (iv) above; or is selected from:
 - (v) OSO_3H , PO_3H_2 , or OPO_3H_2 ;
 - (vi) Y as defined above, but excluding guanidine and quarternary amine salts;
 - (vii) $\text{NHCO}(\text{CH}_2)_m[\text{NHCO}(\text{CH}_2)_{m'}]_p[\text{NHCO}(\text{CH}_2)_{m''}]_q\text{Y}$ where p and q are each 0 or 1, and m, m' and m'' are each an integer from 1 to 10; and
 - 25 (viii) NHCOR^{12} or $\text{NHSO}_2\text{R}^{13}$, where R^{12} and R^{13} are each alkyl groups optionally comprising one or more heteroatoms, and which are substituted by one or more substituents selected from OH, NH_2 , halogen and NO_2 ;
 - (ix) an ether or polyether optionally substituted by one or more hydroxyl groups;
 with the proviso that the compound is other than:

12

- N-{4-Methyl-5-[2-(3-nitro-phenylamino)-pyrimidin-4-yl]-thiazol-2-yl}-methanesulfonamide;
2-Chloro-N-{4-methyl-5-[2-(3-nitro-phenylamino)-pyrimidin-4-yl]-thiazol-2-yl}-acetamide; or
5 2-Chloro-N-{5-[2-(4-fluoro-phenylamino)-pyrimidin-4-yl]-4-methyl-thiazol-2-yl}-acetamide.

- Preferably, the compounds of formula I bear a mono- or di-substituted thiazol-3-yl or thiazol-5-yl radical attached to the pyrimidine ring through one of the ring carbon atoms
10 Most preferably, the heterocycle is a thiazol-5-yl group.

Thus, in one preferred embodiment of the invention, X¹ is S and X² is N.

- The following preferred features apply to compounds of formula I, Ia, Ib and Ic.
15

In another preferred embodiment, Z is NH.

In another preferred embodiment, R³ is H.

- 20 In yet another preferred embodiment, at least one of R², R⁵, R⁶ or R⁷ is an R⁹ or R¹⁰-containing group, or is R¹¹.

- In one particularly preferred embodiment, X¹ is S, X² is N, Z is NH, R¹ is Me, R² is alkyl or amino, R³ is H, one or two of R⁵, R⁶, and R⁷ are CF₃, OH, O-alkyl, halogeno, NO₂, NH₂,
25 NH-alkyl or N-(alkyl)₂ and at least one of R², R⁵, R⁶ or R⁷ is an R⁹ or R¹⁰-containing group, or is R¹¹.

In another preferred embodiment, at least one of R¹, R², R³, R⁴, R⁵, R⁶, R⁷ and R⁸ is R¹¹.

In one preferred embodiment, R^{11} is a solubilising group as defined for R^9 and R^{10} in (i)-(iv) above, or (v)-(x) as defined above.

In another preferred embodiment, R^{11} is a solubilising group as defined for R^9 and R^{10} in (i)-(iv) above, or (v)-(vii), (ix)-(x) as defined above, or is selected from:

- $(CH_2)_nNR^{14}COR^{12}$, where R^{12} is an alkyl group optionally comprising one or more heteroatoms, and which is optionally substituted by one or more substituents selected from OH, NH_2 and NO_2 ,
 - $(CH_2)_nNR^{15}SO_2R^{13}$, where R^{13} is an alkyl group optionally comprising one or more heteroatoms, and which is substituted by one or more substituents selected from OH, NH_2 , halogen and NO_2 ,
 - SO_2R^{16} , where R^{16} is an alkyl group optionally comprising one or more heteroatoms, and which is optionally substituted by one or more substituents selected from OH, NH_2 , halogen and NO_2 ; and
- R^{14} and R^{15} are each independently H or alkyl, and n and n' are each independently 0, 1, 2, or 3.

Preferably, the solubilising group is R^{11} and is:

- (a) Y as defined in above, but excluding guanidine, where Y can also be an alicyclic, aromatic, or heterocyclic group comprising one or more =N- groups;
- (b) $NHCO(CH_2)_m[NHCO(CH_2)_{m'}]_p[NHCO(CH_2)_{m''}]_qY$ or $NHCO(CH_2)_tNH(CH_2)_{t'}Y$ where p and q are each 0 or 1, and m, m', m'', t and t' are each an integer from 1 to 10; or
- (c) $(CH_2)_nNR^{14}COR^{12}$, $(CH_2)_nNR^{15}SO_2R^{13}$, or SO_2R^{16} , where R^{12} , R^{13} and R^{16} are each alkyl groups optionally comprising one or more heteroatoms, and which are substituted by one or more substituents selected from OH, NH_2 , halogen and NO_2 , R^{14} and R^{15} are each independently H or alkyl, and n and n' are each independently 0, 1, 2, or 3.

Preferably, the solubilising group is R^{11} , and R^{11} is:

14

- (a) Y as defined above, but excluding guanidine, where Y can also be an alicyclic, aromatic, or heterocyclic group comprising one or more =N- groups;
- (b) $\text{NHCO}(\text{CH}_2)_m[\text{NHCO}(\text{CH}_2)_{m'}]_p[\text{NHCO}(\text{CH}_2)_{m''}]_q\text{Y}$ where p and q are each 0 or 1, and m, m' and m'' are each integers from 1 to 10
- 5 (c) NHCOR^{12} or $\text{NHSO}_2\text{R}^{13}$, where R^{12} and R^{13} are each alkyl groups optionally comprising one or more heteroatoms, and which are optionally substituted by one or more substituents selected from OH, NH_2 , halogen and NO_2 .

Even more preferably, Y is an alicyclic group comprising one or more of the functions -O-,
 10 NH_2 , -NH-, =N-, a quarternary amine salt, or amidine, and wherein Y is optionally substituted by one or more substituents as defined above.























More preferably still, Y is a morpholine or piperazine group, each of which may be optionally substituted by one or more substituents selected from SO_2 -alkyl, alkyl
 15 optionally substituted by one or more OH groups, CO-alkyl, aralkyl, COO-alkyl, and an ether group optionally substituted by one or more OH groups


In one especially preferred embodiment of the invention, Y is a 2-oxo-hexahydro-thien[3,4-d]imidazole group.

20 In one preferred embodiment, at least one of R^2 , R^6 or R^7 is R^{11} .

For this embodiment, preferably R^{11} is selected from the following:


15






- In one especially preferred embodiment, R^6 or R^7 is R^{11} . More preferably, R^6 is R^{11} and R^2 , R^4 , R^5 , R^7 and R^8 are each independently selected from alkyl, H, CF_3 , OH, O-alkyl, halogeno, NO_2 , NH_2 , NH-alkyl and N-(alkyl) $_2$. More preferably still, R^6 is R^{11} and R^2 , R^4 , R^5 , R^7 and R^8 are each independently selected from alkyl, H, O-alkyl, halogeno, NO_2 , NH_2 and NH-alkyl. Even more preferably, R^6 is R^{11} and R^4 , R^5 , R^7 and R^8 are all H and R^2 is selected from alkyl, O-alkyl, NH_2 and NH-alkyl.

- 10 Even more preferably still, for this embodiment, R^{11} is selected from:



17


In another preferred embodiment, R^7 is R^{11} and R^4, R^5, R^6, R^8 are all H, and R^2 is selected from alkyl, O-alkyl, NH_2 and NH-alkyl. Preferably, for this embodiment, R^{11} is selected from:




5

In another preferred embodiment of the invention, at least one of R^2 or R^6 is R^{11} .

For this embodiment, R^{11} is preferably selected from the following:



10



In one especially preferred embodiment, R^6 is R^{11} .


- 5 For this embodiment, where R^6 is R^{11} , preferably R^2 , R^4 , R^5 , R^7 and R^8 are each independently selected from alkyl, H, CF_3 , OH, O-alkyl, halogeno, NO_2 , NH_2 , NH-alkyl and N-(alkyl)₂.

- Even more preferably, R^2 , R^4 , R^5 , R^7 and R^8 are each independently selected from alkyl, H,
 10 O-alkyl, halogeno, NO_2 , NH_2 and NH-alkyl.

19

More preferably still, R^4 , R^5 , R^7 and R^8 are all H and R^2 is selected from alkyl, O-alkyl, NH_2 and NH-alkyl.

5 More preferably still, R^{11} is selected from:



In an alternative preferred embodiment, R^2 is R^{11} .


10

For this embodiment, R^2 is R^{11} , preferably R^4 , R^5 , R^6 , R^7 and R^8 are each independently selected from alkyl, H, CF_3 , OH, O-alkyl, halogeno, NO_2 , NH_2 , NH-alkyl and N-(alkyl)₂.

15 More preferably, R^4 , R^5 , R^6 , R^7 and R^8 are each independently selected from H, O-alkyl, halogeno, N-(alkyl)₂, NO_2 .


More preferably still, one of R^5 or R^7 is selected from NO_2 , alkoxy, halogeno and N-(alkyl)₂, and the remainder of R^4 , R^5 , R^6 , R^7 and R^8 are all H.

20 More preferably still, R^{11} is selected from:



In one preferred embodiment of the invention, R^1 is methyl, Z is NH and R^3 is H.

- 5 In another embodiment, the compound of the invention is of formula Id, or or a pharmaceutically acceptable salt thereof,



Id

- Preferably, R^1 and R^{3-8} are as defined above for compounds of formula I, Ia, Ib and Ic.
 5 Preferably, R^{17} is alkyl, more preferably methyl.

In a preferred embodiment of the invention, the compound of formula I is selected from those listed in Table 1, but excluding compounds [24], [32] and [33].

- 10 In one especially preferred embodiment, the compound of formula I is selected from the following:
- [4-(2-Methoxy-4-methyl-thiazol-5-yl)-pyrimidin-2-yl]-(4-morpholin-4-yl-phenyl)-amine;
 - [4-(2-Amino-4-methyl-thiazol-5-yl)-pyrimidin-2-yl]-(4-morpholin-4-yl-phenyl)-amine;
 - [4-(2,4-Dimethyl-thiazol-5-yl)-pyrimidin-2-yl]-(4-morpholin-4-yl-phenyl)-amine;
 - 15 [4-(2-N-Methylamino-4-methyl-thiazol-5-yl)-pyrimidin-2-yl]-(4-morpholinophenyl)-amine;
 - [4-(2-Ethylamino-4-methyl-thiazol-5-yl)-pyrimidin-2-yl]-(4-morpholin-4-yl-phenyl)-amine;
 - 1-(4-{4-[4-(2,4-Dimethyl-thiazol-5-yl)-pyrimidin-2-ylamino]-phenyl}-piperazin-1-yl)-ethanone;
 - 20 [4-(2,4-Dimethyl-thiazol-5-yl)-pyrimidin-2-yl]-(4-piperazin-1-yl-phenyl)-amine;
 - [4-(2,4-Dimethyl-thiazol-5-yl)-pyrimidin-2-yl]-[4-(4'-2"-ethoxylethanolpiperazino)-phenyl]-amine;
 - 3-(4-{4-[4-(2,4-Dimethyl-thiazol-5-yl)-pyrimidin-2-ylamino]-phenyl}-piperazin-1-yl)-propan-1-ol;
 - 25

- 2-(4-{4-[4-(2,4-Dimethyl-thiazol-5-yl)-pyrimidin-2-ylamino]-phenyl}-piperazin-1-yl)-ethanol;
- [4-(2,4-Dimethyl-thiazol-5-yl)-pyrimidin-2-yl]-[4-(4-methanesulfonyl-piperazin-1-yl)-phenyl]-amine;
- 5 [4-(4-Benzyl-piperazin-1-yl)-phenyl]-[4-(4-methyl-2-methylamino-thiazol-5-yl)-pyrimidin-2-yl]-amine;
- [4-(2,4-Dimethyl-thiazol-5-yl)-pyrimidin-2-yl]-[4-(4-methyl-piperazin-1-yl)-phenyl]-amine;
- [4-(2-Methoxy-4-methyl-thiazol-5-yl)-pyrimidin-2-yl]-[4-(4-methyl-piperazin-1-yl)-phenyl]-amine;
- 10 3-Amino-N-{4-methyl-5-[2-(3-nitro-phenylamino)-pyrimidin-4-yl]-thiazol-2-yl}-propionamide;
- (2S)-2-Amino-3-hydroxy-N-{4-methyl-5-[2-(3-nitro-phenylamino)-pyrimidin-4-yl]-thiazol-2-yl}-propionamide;
- 15 (2R,3R)-2-Amino-3-hydroxy-N-{4-methyl-5-[2-(3-nitro-phenylamino)-pyrimidin-4-yl]-thiazol-2-yl}-butyramide;
- (2R)-2-Amino-N-{4-methyl-5-[2-(3-nitro-phenylamino)-pyrimidin-4-yl]-thiazol-2-yl}-butyramide;
- (2S,3S)-2-Amino-3-hydroxy-N-{4-methyl-5-[2-(3-nitro-phenylamino)-pyrimidin-4-yl]-thiazol-2-yl}-butyramide;
- 20 4-Amino-N-{4-methyl-5-[2-(3-nitro-phenylamino)-pyrimidin-4-yl]-thiazol-2-yl}-butyramide;
- 3-Amino-N-{4-methyl-5-[2-(4-morpholin-4-yl-phenylamino)-pyrimidin-4-yl]-thiazol-2-yl}-propionamide;
- 25 3-Bromo-N-{5-[2-(4-fluoro-phenylamino)-pyrimidin-4-yl]-4-methyl-thiazol-2-yl}-propionamide;
- N-{5-[2-(4-Fluoro-phenylamino)-pyrimidin-4-yl]-4-methyl-thiazol-2-yl}-3-morpholin-4-yl-propionamide;
- N-{4-Methyl-5-[2-(3-nitro-phenylamino)-pyrimidin-4-yl]-thiazol-2-yl}-3-morpholin-4-yl-propionamide;
- 30

- N-{4-Methyl-5-[2-(3-nitro-phenylamino)-pyrimidin-4-yl]-thiazol-2-yl}-3-(4-methyl-piperazin-1-yl)-propionamide;
- 2-Chloro-N-{5-[2-(3-chloro-phenylamino)-pyrimidin-4-yl]-4-methyl-thiazol-2-yl}-acetamide;
- 5 2-Chloro-N-{5-[2-(3-methoxy-phenylamino)-pyrimidin-4-yl]-4-methyl-thiazol-2-yl}-acetamide;
- 2-Chloro-N-{5-[2-(4-dimethylamino-phenylamino)-pyrimidin-4-yl]-4-methyl-thiazol-2-yl}-acetamide;
- 4-({4-Methyl-5-[2-(3-nitro-phenylamino)-pyrimidin-4-yl]-thiazol-2-ylcarbamoyl}-methyl)-piperazine-1-carboxylic acid tert-butyl ester;
- 10 N-{5-[2-(4-Dimethylamino-phenylamino)-pyrimidin-4-yl]-4-methyl-thiazol-2-yl}-2-[2-(2-hydroxy-ethoxy)-ethylamino]-acetamide;
- 6-[5-(2-Oxo-hexahydro-thieno[3,4-d]imidazol-4-yl)-pentanoylamino]-hexanoic acid (2-{4-methyl-5-[2-(4-morpholin-4-yl-phenylamino)-pyrimidin-4-yl]-thiazol-2-ylcarbamoyl}-ethyl)-amide;
- 15 N-{5-[2-(4-Fluoro-phenylamino)-pyrimidin-4-yl]-4-methyl-thiazol-2-yl}-methanesulfonamide;
- 3-Bromo-N-{4-methyl-5-[2-(3-nitro-phenylamino)-pyrimidin-4-yl]-thiazol-2-yl}-propionamide;
- 20 3-(1-{4-[4-(2,4-Dimethyl-thiazol-5-yl)-pyrimidin-2-ylamino]-phenyl}-piperidin-4-yl)-propan-1-ol;
- 2-Amino-3-hydroxy-N-{4-methyl-5-[2-(3-nitro-phenylamino)-pyrimidin-4-yl]-thiazol-2-yl}-butyramide; and
- 2-Chloro-N-{5-[2-(4-chloro-phenylamino)-pyrimidin-4-yl]-4-methyl-thiazol-2-yl}-acetamide.
- 25

In a further preferred embodiment of the invention, said compound of formula I is selected from the following:

- [4-(2-Amino-4-methyl-thiazol-5-yl)-pyrimidin-2-yl]-(4-morpholin-4-yl-phenyl)-amine;
- 30 [4-(2,4-dimethyl-thiazol-5-yl)-pyrimidin-2-yl]-(4-morpholin-4-yl-phenyl)-amine;

[4-(2-*N*-methylamino-4-methyl-thiazol-5-yl)-pyrimidin-2-yl]-(4-morpholinophenyl)-amine;

(2*R*,3*R*)-2-amino-3-hydroxy-*N*-{4-methyl-5-[2-(3-nitro-phenylamino)-pyrimidin-4-yl]-thiazol-2-yl}-butyramide;

5 (2*R*)-2-amino-*N*-{4-methyl-5-[2-(3-nitro-phenylamino)-pyrimidin-4-yl]-thiazol-2-yl}-butyramide; and

N-{4-methyl-5-[2-(3-nitro-phenylamino)-pyrimidin-4-yl]-thiazol-2-yl}-3-morpholin-4-yl-propionamide.

10 In one preferred embodiment the compound of formula I is capable of exhibiting an antiproliferative effect in human cell lines, as measured by a standard 72h MTT cytotoxicity assay. Preferably, the compound of formula I exhibits an IC₅₀ value of less than 10 μ M, more preferably less than 5 μ M, even more preferably less than 1 μ M as measured by said MTT assay. More preferably, the compound of formula I is selected from
15 the following: [4], [8], [12], [14], [16], [22], [24], [25], [29], [32], [33], [39], [40], [50], [53], [61], [57], [62], [63], [64], [65] and [77]. More preferably still, the compound exhibits an IC₅₀ value of less than 0.5 μ M, more preferably still less than 0.2 μ M. Even more preferably, the compound is selected from the following: [24], [25], [32], [33], [50], [62] and [64].

20

In another preferred embodiment, the compound of formula I is selected from [1], [11], [15] and [16].

In another preferred embodiment, the compound of formula I is capable of inhibiting one
25 or more protein kinases, as measured by the assays described in the accompanying Examples section. Preferably, the compound of formula I exhibits an IC₅₀ value of less than 10 μ M, more preferably less than 5 μ M, even more preferably less than 1 μ M or less than 0.5 μ M, more preferably still less than 0.1 μ M. More preferably, the compound of formula I is selected from the following: [11], [13], [14], [15], [20], [21], [22], [24],
30 [25], [32], [50], [53], [54], [55], [56], [57], [59], [61], [62], [64], [68], [71], [82], [83], [84]

25

and [85]. More preferably still, the compound exhibits an IC_{50} value of less than $0.01 \mu M$. Even more preferably, the compound is selected from the following: [11], [22], [24], [32], [50], [62], [71] and [85].

- 5 In yet another preferred embodiment, the compound of formula I is selected from [1], [3], [11], [15] and [16].

The present invention provides a series of compounds equipped with solubilising functions on the phenyl and/or heteroaryl rings of the 2-phenylamino-4-heteroaryl-pyrimidine
10 system. Modification with solubilising moieties has preserved the desired *in vitro* biological activity (inhibition of CDKs and cytotoxicity against transformed human cells) and in some cases has led to surprising and unexpected increases in potency. Furthermore, *in vivo* absorption, and oral bioavailability in particular can also be improved using the solubilising strategies presented herein.

15

THERAPEUTIC USE

The compounds of formula I have been found to possess anti-proliferative activity and are therefore believed to be of use in the treatment of proliferative disorders such as cancers, leukaemias and other disorders associated with uncontrolled cellular proliferation such as
20 psoriasis and restenosis. As defined herein, an anti-proliferative effect within the scope of the present invention may be demonstrated by the ability to inhibit cell proliferation in an *in vitro* whole cell assay, for example using any of the cell lines AGS, H1299 or SJSA-1, or by showing inhibition of the interaction between HDM2 and p53 in an appropriate assay. These assays, including methods for their performance, are described in more detail
25 in the accompanying Examples. Using such assays it may be determined whether a compound is anti-proliferative in the context of the present invention.

On preferred embodiment of the present invention therefore relates to the use of one or more compounds of formula I in the treatment of proliferative disorders. Preferably, the
30 proliferative disorder is a cancer or leukaemia. The term proliferative disorder is used

herein in a broad sense to include any disorder that requires control of the cell cycle, for example cardiovascular disorders such as restenosis and cardiomyopathy, auto-immune disorders such as glomerulonephritis and rheumatoid arthritis, dermatological disorders such as psoriasis, anti-inflammatory, anti-fungal, antiparasitic disorders such as malaria, emphysema and alopecia. In these disorders, the compounds of the present invention may induce apoptosis or maintain stasis within the desired cells as required.

The compounds of the invention may inhibit any of the steps or stages in the cell cycle, for example, formation of the nuclear envelope, exit from the quiescent phase of the cell cycle (G0), G1 progression, chromosome decondensation, nuclear envelope breakdown, START, initiation of DNA replication, progression of DNA replication, termination of DNA replication, centrosome duplication, G2 progression, activation of mitotic or meiotic functions, chromosome condensation, centrosome separation, microtubule nucleation, spindle formation and function, interactions with microtubule motor proteins, chromatid separation and segregation, inactivation of mitotic functions, formation of contractile ring, and cytokinesis functions. In particular, the compounds of the invention may influence certain gene functions such as chromatin binding, formation of replication complexes, replication licensing, phosphorylation or other secondary modification activity, proteolytic degradation, microtubule binding, actin binding, septin binding, microtubule organising centre nucleation activity and binding to components of cell cycle signalling pathways.

In one embodiment of the invention, the compound of formula I or Ia is administered in an amount sufficient to inhibit at least one CDK enzyme.


In a more preferred embodiment of the invention, the compound of formula Ia is preferably administered in an amount sufficient to inhibit one or more of the host cell CDKs involved in viral replication, *i.e.* CDK2, CDK7, CDK8, and CDK9 [Wang D, De la Fuente C, Deng L, Wang L, Zilberman I, Eadie C, Healey M, Stein D, Denny T, Harrison LE, Meijer L, Kashanchi F. Inhibition of human immunodeficiency virus type 1 transcription by chemical cyclin-dependent kinase inhibitors. *J. Virol.* 2001; 75: 7266-7279].

As defined herein, an anti-viral effect within the scope of the present invention may be demonstrated by the ability to inhibit CDK2, CDK7, CDK8 or CDK9. Assays for determining CDK activity are described in more detail in the accompanying examples. Using such enzymes assays it may be determined whether a compound is anti-viral in the context of the present invention.

In a particularly preferred embodiment, the compounds of formula Ia are useful in the treatment of viral disorders, such as human cytomegalovirus (HCMV), herpes simplex virus type 1 (HSV-1), human immunodeficiency virus type 1 (HIV-1), and varicella zoster virus (VZV).

In a particularly preferred embodiment, the invention relates to the use of one or more compounds of formula Ia in the treatment of a viral disorder which is CDK dependent or sensitive. CDK dependent disorders are associated with an above normal level of activity of one or more CDK enzymes. Such disorders preferably associated with an abnormal level of activity of CDK2, CDK7, CDK8 and/or CDK9. A CDK sensitive disorder is a disorder in which an aberration in the CDK level is not the primary cause, but is downstream of the primary metabolic aberration. In such scenarios, CDK2, CDK7, CDK8 and/or CDK9 can be said to be part of the sensitive metabolic pathway and CDK inhibitors may therefore be active in treating such disorders.

One preferred embodiment of the invention relates to the use of a compound of formula Ie, or a pharmaceutically acceptable salt thereof,



Ie

wherein

one of X^1 and X^2 is S, and the other of X^1 and X^2 is N;

Z is NH, NHCO, NHSO₂, NHCH₂, CH₂, CH₂CH₂, CH=CH, SO₂, or SO;

5

R^1 , R^2 , R^3 , R^4 , R^5 , R^6 , R^7 and R^8 are each independently H, alkyl, alkyl- R^9 , aryl, aryl- R^9 , aralkyl, aralkyl- R^9 , halogeno, NO₂, CN, OH, O-alkyl, COR⁹, COOR⁹, O-aryl, O- R^9 , NH₂, NH-alkyl, NH-aryl, N-(alkyl)₂, N-(aryl)₂, N-(alkyl)(aryl), NH- R^9 , N-(R^9)(R^{10}), N-(alkyl)(R^9), N-(aryl)(R^9), COOH, CONH₂, CONH-alkyl, CONH-aryl, CON-(alkyl)(R^9),
 10 CON(aryl)(R^9), CONH- R^9 , CON-(R^9)(R^{10}), SO₃H, SO₂-alkyl, SO₂-alkyl- R^9 , SO₂-aryl, SO₂-aryl- R^9 , SO₂NH₂, SO₂NH- R^9 , SO₂N-(R^9)(R^{10}), CF₃, CO-alkyl, CO-alkyl- R^9 , CO-aryl, CO-aryl- R^9 or R^{11} , wherein alkyl, aryl, aralkyl groups may be further substituted with one or more groups selected from halogeno, NO₂, OH, O-methyl, NH₂, COOH, CONH₂ and CF₃;

15

wherein at least one of R^1 , R^2 , R^3 , R^4 , R^5 , R^6 , R^7 and R^8 is an R^9 or R^{10} -containing group, or is R^{11} ;

R^9 and R^{10} are each independently solubilising groups selected from:

- 20 (i) - a mono-, di- or polyhydroxylated alicyclic group;
 - a di- or polyhydroxylated aliphatic or aromatic group;
 - a carbohydrate derivative;
 - an O- and/or S-containing heterocyclic group optionally substituted by one or more hydroxyl groups;
 25 - an aliphatic or aromatic group containing a carboxamide, sulfoxide, sulfone, or sulfonamide function; or
 - a halogenated alkylcarbonyl group;
- (ii) COOH, SO₃H, OSO₃H, PO₃H₂, or OPO₃H₂;
- (iii) Y, where Y is selected from an alicyclic, aromatic, or heterocyclic group
 30 comprising one or more of the functions =N-, -O-, -NH₂, -NH-, a quaternary amine

29

salt, guanidine, and amidine, where Y is optionally substituted by one or more substituents selected from:

- SO₂-alkyl;
- alkyl optionally substituted by one or more OH groups;
- 5 - CO-alkyl;
- aralkyl;
- COO-alkyl; and
- an ether group optionally substituted by one or more OH groups; and

where Y is other than pyridinyl;

- 10 (iv) a natural or unnatural amino acid, a peptide or a peptide derivative;

R¹¹ is a solubilising group as defined for R⁹ and R¹⁰ in (i) or (iv) above; or is selected from:

- (v) OSO₃H, PO₃H₂, or OPO₃H₂;
 - (vi) Y as defined above, but excluding guanidine and quaternary amine salts;
 - 15 (vii) NHCO(CH₂)_m[NHCO(CH₂)_{m'}]_p[NHCO(CH₂)_{m''}]_qY where p and q are each 0 or 1, and m, m' and m'' are each an integer from 1 to 10; and
 - (viii) NHCOR¹² or NHSO₂R¹³, where R¹² and R¹³ are each alkyl groups optionally comprising one or more heteroatoms, and which are substituted by one or more substituents selected from OH, NH₂, halogen and NO₂;
 - 20 (ix) an ether or polyether optionally substituted by one or more hydroxyl groups;
- in the preparation of a medicament for treating a viral disorder.

Preferred features are as defined above for compounds of formula I, Ia, Ib and Ic.

- 25 In a preferred embodiment, the compound of formula Ia is selected from those listed in Table 1.

More preferably, said compound of formula Ia is selected from the following: [1], [3], [4], [15] and [53].

For use in the treatment of viral disorders, preferably the compound of formula Ia is capable of inhibiting CK2, CDK7 and/or CDK9 and is selected from the following:

[4-(2-Ethylamino-4-methyl-thiazol-5-yl)-pyrimidin-2-yl]-(4-morpholin-4-yl-phenyl)-amine;

2-Amino-3-hydroxy-N-{4-methyl-5-[2-(3-nitro-phenylamino)-pyrimidin-4-yl]-thiazol-2-yl}-butyramide;

2-Amino-N-{4-methyl-5-[2-(3-nitro-phenylamino)-pyrimidin-4-yl]-thiazol-2-yl}-butyramide;

[4-(2-Amino-4-methyl-thiazol-5-yl)-pyrimidin-2-yl]-(4-morpholin-4-yl-phenyl)-amine;

[4-(4-Methyl-2-methylamino-thiazol-5-yl)-pyrimidin-2-yl]-(4-morpholin-4-yl-phenyl)-amine;

3-(1-{4-[4-(2,4-Dimethyl-thiazol-5-yl)-pyrimidin-2-ylamino]-phenyl}-piperidin-4-yl)-propan-1-ol;

N-{5-[2-(4-Fluoro-phenylamino)-pyrimidin-4-yl]-4-methyl-thiazol-2-yl}-3-morpholin-4-yl-propionamide;

3-Bromo-N-{4-methyl-5-[2-(3-nitro-phenylamino)-pyrimidin-4-yl]-thiazol-2-yl}-propionamide;

N-{4-Methyl-5-[2-(3-nitro-phenylamino)-pyrimidin-4-yl]-thiazol-2-yl}-3-morpholin-4-yl-propionamide;

N-{4-Methyl-5-[2-(3-nitro-phenylamino)-pyrimidin-4-yl]-thiazol-2-yl}-3-(4-methyl-piperazin-1-yl)-propionamide;

[4-(2-Methoxy-4-methyl-thiazol-5-yl)-pyrimidin-2-yl]-[4-(4-methyl-piperazin-1-yl)-phenyl]-amine;

[4-(4-Benzyl-piperazin-1-yl)-phenyl]-[4-(4-methyl-2-methylamino-thiazol-5-yl)-pyrimidin-2-yl]-amine; and

3-Amino-N-{4-methyl-5-[2-(3-nitro-phenylamino)-pyrimidin-4-yl]-thiazol-2-yl}-propionamide.

The following compound is observed to be a particularly effective anti-viral agent, as demonstrated by cell based assays: [4-(2-Amino-4-methyl-thiazol-5-yl)-pyrimidin-2-yl]-(4-morpholin-4-yl-phenyl)-amine.

- 5 Another aspect of the invention relates to the use of a compound of formula I or Ia as an anti-mitotic agent.

Yet another aspect of the invention relates to the use of a compound of formula I or Ia for treating a neurodegenerative disorder.

10

Preferably, the neurodegenerative disorder is neuronal apoptosis.

Another aspect of the invention relates to the use of a compound of formula I or Ia as an antiviral agent.

15

Thus, another aspect of the invention relates to the use of a compound of the invention in the preparation of a medicament for treating a viral disorder, such as human cytomegalovirus (HCMV), herpes simplex virus type 1 (HSV-1), human immunodeficiency virus type 1 (HIV-1), and varicella zoster virus (VZV).

20

In a more preferred embodiment of the invention, the compound of the invention is administered in an amount sufficient to inhibit one or more of the host cell CDKs involved in viral replication, i.e. CDK2, CDK7, CDK8, and CDK9 [Wang D, De la Fuente C, Deng L, Wang L, Zilberman I, Eadie C, Healey M, Stein D, Denny T, Harrison LE, Meijer L, Kashanchi F. Inhibition of human immunodeficiency virus type 1 transcription by chemical cyclin-dependent kinase inhibitors. J. Virol. 2001; 75: 7266-7279].

25

As defined herein, an anti-viral effect within the scope of the present invention may be demonstrated by the ability to inhibit CDK2, CDK7, CDK8 or CDK9.

30

In a particularly preferred embodiment, the invention relates to the use of one or more compounds of the invention in the treatment of a viral disorder which is CDK dependent or sensitive. CDK dependent disorders are associated with an above normal level of activity of one or more CDK enzymes. Such disorders preferably associated with an abnormal
5 level of activity of CDK2, CDK7, CDK8 and/or CDK9. A CDK sensitive disorder is a disorder in which an aberration in the CDK level is not the primary cause, but is downstream of the primary metabolic aberration. In such scenarios, CDK2, CDK7, CDK8 and/or CDK9 can be said to be part of the sensitive metabolic pathway and CDK inhibitors may therefore be active in treating such disorders.

10

Another aspect of the invention relates to the use of compounds of formula I or Ia, or pharmaceutically acceptable salts thereof, in the preparation of a medicament for treating diabetes.

15 In a particularly preferred embodiment, the diabetes is type II diabetes.

GSK3 is one of several protein kinases that phosphorylate glycogen synthase (GS). The stimulation of glycogen synthesis by insulin in skeletal muscle results from the dephosphorylation and activation of GS. GSK3's action on GS thus results in the latter's
20 deactivation and thus suppression of the conversion of glucose into glycogen in muscles.

Type II diabetes (non-insulin dependent diabetes mellitus) is a multi-factorial disease. Hyperglycaemia is due to insulin resistance in the liver, muscles, and other tissues, coupled with impaired secretion of insulin. Skeletal muscle is the main site for insulin-stimulated
25 glucose uptake, there it is either removed from circulation or converted to glycogen. Muscle glycogen deposition is the main determinant in glucose homeostasis and type II diabetics have defective muscle glycogen storage. There is evidence that an increase in GSK3 activity is important in type II diabetes [Chen, Y.H.; Hansen, L.; Chen, M.X.; Bjorbaek, C.; Vestergaard, H.; Hansen, T.; Cohen, P.T.; Pedersen, O. *Diabetes*, 1994, 43,
30 1234]. Furthermore, it has been demonstrated that GSK3 is over-expressed in muscle cells

of type II diabetics and that an inverse correlation exists between skeletal muscle GSK3 activity and insulin action [Nikoulina, S.E.; Ciaraldi, T.P.; Mudaliar, S.; Mohideen, P.; Carter, L.; Henry, R.R. *Diabetes*, 2000, 49, 263].

- 5 GSK3 inhibition is therefore of therapeutic significance in the treatment of diabetes, particularly type II, and diabetic neuropathy.

It is notable that GSK3 is known to phosphorylate many substrates other than GS, and is thus involved in the regulation of multiple biochemical pathways. For example, GSK is
10 highly expressed in the central and peripheral nervous systems.

Another aspect of the invention therefore relates to the use of compounds of formula I or Ia, or pharmaceutically acceptable salts thereof, in the preparation of a medicament for treating a CNS disorders, for example neurodegenerative disorders.

15

Preferably, the CNS disorder is Alzheimer's disease.

Tau is a GSK-3 substrate which has been implicated in the etiology of Alzheimer's disease. In healthy nerve cells, Tau co-assembles with tubulin into microtubules. However, in
20 Alzheimer's disease, tau forms large tangles of filaments, which disrupt the microtubule structures in the nerve cell, thereby impairing the transport of nutrients as well as the transmission of neuronal messages.

Without wishing to be bound by theory, it is believed that GSK3 inhibitors may be able to
25 prevent and/or reverse the abnormal hyperphosphorylation of the microtubule-associated protein tau that is an invariant feature of Alzheimer's disease and a number of other neurodegenerative diseases, such as progressive supranuclear palsy, corticobasal degeneration and Pick's disease. Mutations in the tau gene cause inherited forms of fronto-temporal dementia, further underscoring the relevance of tau protein dysfunction for the
30 neurodegenerative process [Goedert, M. *Curr. Opin. Gen. Dev.*, 2001, 11, 343].

Another aspect of the invention relates to the use of compounds of formula I or Ia, or pharmaceutically acceptable salts thereof, in the preparation of a medicament for treating bipolar disorder.

- 5 Yet another aspect of the invention relates to the use of compounds of formula I or Ia, or pharmaceutically acceptable salts thereof, in the preparation of a medicament for treating a stroke.

Reducing neuronal apoptosis is an important therapeutic goal in the context of head
10 trauma, stroke, epilepsy, and motor neuron disease [Mattson, M.P. Nat. Rev. Mol. Cell. Biol., 2000, 1, 120]. Therefore, GSK3 as a pro-apoptotic factor in neuronal cells makes this protein kinase an attractive therapeutic target for the design of inhibitory drugs to treat these diseases.

- 15 Yet another aspect of the invention relates to the use of compounds of formula I or Ia, or pharmaceutically acceptable salts thereof, in the preparation of a medicament for treating alopecia.

Hair growth is controlled by the Wnt signalling pathway, in particular Wnt-3. In tissue-
20 culture model systems of the skin, the expression of non-degradable mutants of β -catenin leads to a dramatic increase in the population of putative stem cells, which have greater proliferative potential [Zhu, A.J.; Watt, F.M. Development, 1999, 126, 2285]. This population of stem cells expresses a higher level of non-cadherin-associated β -catenin [DasGupta, R.; Fuchs, E. Development, 1999, 126, 4557], which may contribute to their
25 high proliferative potential. Moreover, transgenic mice overexpressing a truncated β -catenin in the skin undergo de novo hair-follicle morphogenesis, which normally is only established during embryogenesis. The ectopic application of GSK3 inhibitors may therefore be therapeutically useful in the treatment of baldness and in restoring hair growth following chemotherapy-induced alopecia.

A further aspect of the invention relates to a method of treating a GSK3-dependent disorder, said method comprising administering to a subject in need thereof, a compound of formula I or Ia, or a pharmaceutically acceptable salt thereof, as defined above in an amount sufficient to inhibit GSK3.

5

Preferably, the compound of the invention, or pharmaceutically acceptable salt thereof, is administered in an amount sufficient to inhibit GSK3 β .

In one embodiment of the invention, the compound of the invention is administered in an amount sufficient to inhibit at least one PLK enzyme.

10

The polo-like kinases (PLKs) constitute a family of serine/threonine protein kinases. Mitotic *Drosophila melanogaster* mutants at the polo locus display spindle abnormalities [Sunkel et al., *J. Cell Sci.*, 1988, 89, 25] and polo was found to encode a mitotic kinase [Llamazares et al., *Genes Dev.*, 1991, 5, 2153]. In humans, there exist three closely related PLKs [Glover et al., *Genes Dev.*, 1998, 12, 3777]. They contain a highly homologous amino-terminal catalytic kinase domain and their carboxyl termini contain two or three conserved regions, the polo boxes. The function of the polo boxes remains incompletely understood but they are implicated in the targeting of PLKs to subcellular compartments [Lee et al., *Proc. Natl. Acad. Sci. USA*, 1998, 95, 9301; Leung et al., *Nat. Struct. Biol.*, 2002, 9, 719], mediation of interactions with other proteins [Kauselmann et al., *EMBO J.*, 1999, 18, 5528], or may constitute part of an autoregulatory domain [Nigg, *Curr. Opin. Cell Biol.*, 1998, 10, 776]. Furthermore, the polo box-dependent PLK1 activity is required for proper metaphase/anaphase transition and cytokinesis [Yuan et al., *Cancer Res.*, 2002, 62, 4186; Seong et al., *J. Biol. Chem.*, 2002, 277, 32282].

15

20

25

Studies have shown that human PLKs regulate some fundamental aspects of mitosis [Lane et al., *J. Cell. Biol.*, 1996, 135, 1701; Cogswell et al., *Cell Growth Differ.*, 2000, 11, 615]. In particular, PLK1 activity is believed to be necessary for the functional maturation of centrosomes in late G2/early prophase and subsequent establishment of a bipolar spindle.

30

Depletion of cellular PLK1 through the small interfering RNA (siRNA) technique has also confirmed that this protein is required for multiple mitotic processes and completion of cytokinesis [Liu et al., *Proc. Natl. Acad. Sci. USA*, 2002, 99, 8672].

- 5 In a more preferred embodiment of the invention, the compound of the invention is administered in an amount sufficient to inhibit PLK1.

Of the three human PLKs, PLK1 is the best characterized; it regulates a number of cell division cycle effects, including the onset of mitosis [Toyoshima-Morimoto et al., *Nature*,
10 2001, 410, 215; Roshak et al., *Cell. Signalling*, 2000, 12, 405], DNA-damage checkpoint activation [Smits et al., *Nat. Cell Biol.*, 2000, 2, 672; van Vugt et al., *J. Biol. Chem.*, 2001, 276, 41656], regulation of the anaphase promoting complex [Sumara et al., *Mol. Cell*, 2002, 9, 515; Golan et al., *J. Biol. Chem.*, 2002, 277, 15552; Kotani et al., *Mol. Cell*, 1998, 1, 371], phosphorylation of the proteasome [Feng et al., *Cell Growth Differ.*, 2001, 12, 29],
15 and centrosome duplication and maturation [Dai et al., *Oncogene*, 2002, 21, 6195].

Specifically, initiation of mitosis requires activation of M-phase promoting factor (MPF), the complex between the cyclin dependent kinase CDK1 and B-type cyclins [Nurse, *Nature*, 1990, 344, 503]. The latter accumulate during the S and G2 phases of the cell
20 cycle and promote the inhibitory phosphorylation of the MPF complex by WEE1, MIK1, and MYT1 kinases. At the end of the G2 phase, corresponding dephosphorylation by the dual-specificity phosphatase CDC25C triggers the activation of MPF [Nigg, *Nat. Rev. Mol. Cell Biol.*, 2001, 2, 21]. In interphase, cyclin B localizes to the cytoplasm [Hagting et al., *EMBO J.*, 1998, 17, 4127], it then becomes phosphorylated during prophase and this event
25 causes nuclear translocation [Hagting et al., *Curr. Biol.*, 1999, 9, 680; Yang et al., *J. Biol. Chem.*, 2001, 276, 3604]. The nuclear accumulation of active MPF during prophase is thought to be important for initiating M-phase events [Takizawa et al., *Curr. Opin. Cell Biol.*, 2000, 12, 658]. However, nuclear MPF is kept inactive by WEE1 unless counteracted by CDC25C. The phosphatase CDC25C itself, localized to the cytoplasm
30 during interphase, accumulates in the nucleus in prophase [Seki et al., *Mol. Biol. Cell*,

1992, 3, 1373; Heald et al., *Cell*, 1993, 74, 463; Dalal et al., *Mol. Cell. Biol.*, 1999, 19, 4465]. The nuclear entry of both cyclin B [Toyoshima-Morimoto et al., *Nature*, 2001, 410, 215] and CDC25C [Toyoshima-Morimoto et al., *EMBO Rep.*, 2002, 3, 341] are promoted through phosphorylation by PLK1 [Roshak et al., *Cell. Signalling*, 2000, 12, 405]. This
5 kinase is an important regulator of M-phase initiation.

In one particularly preferred embodiment, the compounds of the invention are ATP-antagonistic inhibitors of PLK1.

10 In the present context ATP antagonism refers to the ability of an inhibitor compound to diminish or prevent PLK catalytic activity, i.e. phosphotransfer from ATP to a macromolecular PLK substrate, by virtue of reversibly or irreversibly binding at the enzyme's active site in such a manner as to impair or abolish ATP binding.

15 In another preferred embodiment, the compound of the invention is administered in an amount sufficient to inhibit PLK2 and/or PLK3.

Mammalian PLK2 (also known as SNK) and PLK3 (also known as PRK and FNK) were originally shown to be immediate early gene products. PLK3 kinase activity appears to
20 peak during late S and G2 phase. It is also activated during DNA damage checkpoint activation and severe oxidative stress. PLK3 also plays an important role in the regulation of microtubule dynamics and centrosome function in the cell and deregulated PLK3 expression results in cell cycle arrest and apoptosis [Wang et al., *Mol. Cell. Biol.*, 2002, 22, 3450]. PLK2 is the least well understood homologue of the three PLKs. Both PLK2 and
25 PLK3 may have additional important post-mitotic functions [Kauselmann et al., *EMBO J.*, 1999, 18, 5528].

Another aspect of the invention relates to the use of a compound of formula I for inhibiting a protein kinase.

In a preferred embodiment of this aspect, the protein kinase is a cyclin dependent kinase. Preferably, the protein kinase is CDK1, CDK2, CDK3, CDK4, CDK6, CDK7, CDK8 or CDK9, more preferably CDK2.

- 5 A further aspect of the invention relates to a method of inhibiting a protein kinase, said method comprising contacting said protein kinase with a compound of formula I.

In a preferred embodiment of this aspect, the protein kinase is a cyclin dependent kinase, even more preferably CDK2.

10

PHARMACEUTICAL COMPOSITIONS

- A further aspect of the invention relates to a pharmaceutical composition comprising a compound of formula I as defined for said first aspect admixed with one or more
15 pharmaceutically acceptable diluents, excipients or carriers. Even though the compounds of the present invention (including their pharmaceutically acceptable salts, esters and pharmaceutically acceptable solvates) can be administered alone, they will generally be administered in admixture with a pharmaceutical carrier, excipient or diluent, particularly for human therapy. The pharmaceutical compositions may be for human or animal usage in
20 human and veterinary medicine.

Examples of such suitable excipients for the various different forms of pharmaceutical compositions described herein may be found in the "Handbook of Pharmaceutical Excipients, 2nd Edition, (1994), Edited by A Wade and PJ Weller.

25

Acceptable carriers or diluents for therapeutic use are well known in the pharmaceutical art, and are described, for example, in Remington's Pharmaceutical Sciences, Mack Publishing Co. (A. R. Gennaro edit. 1985).

Examples of suitable carriers include lactose, starch, glucose, methyl cellulose, magnesium stearate, mannitol, sorbitol and the like. Examples of suitable diluents include ethanol, glycerol and water.

- 5 The choice of pharmaceutical carrier, excipient or diluent can be selected with regard to the intended route of administration and standard pharmaceutical practice. The pharmaceutical compositions may comprise as, or in addition to, the carrier, excipient or diluent any suitable binder(s), lubricant(s), suspending agent(s), coating agent(s), solubilising agent(s).

10

Examples of suitable binders include starch, gelatin, natural sugars such as glucose, anhydrous lactose, free-flow lactose, beta-lactose, corn sweeteners, natural and synthetic gums, such as acacia, tragacanth or sodium alginate, carboxymethyl cellulose and polyethylene glycol.

15

Examples of suitable lubricants include sodium oleate, sodium stearate, magnesium stearate, sodium benzoate, sodium acetate, sodium chloride and the like.

- 20 Preservatives, stabilizers, dyes and even flavoring agents may be provided in the pharmaceutical composition. Examples of preservatives include sodium benzoate, sorbic acid and esters of p-hydroxybenzoic acid. Antioxidants and suspending agents may be also used.

SALTS/ESTERS

- 25 The compounds of formula I or Ia can be present as salts or esters, in particular pharmaceutically acceptable salts or esters.

Pharmaceutically acceptable salts of the compounds of the invention include suitable acid addition or base salts thereof. A review of suitable pharmaceutical salts may be found in

Berge et al, J Pharm Sci, 66, 1-19 (1977). Salts are formed, for example with strong inorganic acids such as mineral acids, e.g. sulphuric acid, phosphoric acid or hydrohalic acids; with strong organic carboxylic acids, such as alkanecarboxylic acids of 1 to 4 carbon atoms which are unsubstituted or substituted (e.g., by halogen), such as acetic acid; with
5 saturated or unsaturated dicarboxylic acids, for example oxalic, malonic, succinic, maleic, fumaric, phthalic or tetraphthalic; with hydroxycarboxylic acids, for example ascorbic, glycolic, lactic, malic, tartaric or citric acid; with aminoacids, for example aspartic or glutamic acid; with benzoic acid; or with organic sulfonic acids, such as (C₁-C₄)-alkyl- or aryl-sulfonic acids which are unsubstituted or substituted (for example, by a halogen) such
10 as methane- or p-toluene sulfonic acid.

Esters are formed either using organic acids or alcohols/hydroxides, depending on the functional group being esterified. Organic acids include carboxylic acids, such as alkanecarboxylic acids of 1 to 12 carbon atoms which are unsubstituted or substituted (e.g.,
15 by halogen), such as acetic acid; with saturated or unsaturated dicarboxylic acid, for example oxalic, malonic, succinic, maleic, fumaric, phthalic or tetraphthalic; with hydroxycarboxylic acids, for example ascorbic, glycolic, lactic, malic, tartaric or citric acid; with aminoacids, for example aspartic or glutamic acid; with benzoic acid; or with organic sulfonic acids, such as (C₁-C₄)-alkyl- or aryl-sulfonic acids which are
20 unsubstituted or substituted (for example, by a halogen) such as methane- or p-toluene sulfonic acid. Suitable hydroxides include inorganic hydroxides, such as sodium hydroxide, potassium hydroxide, calcium hydroxide, aluminium hydroxide. Alcohols include alkanealcohols of 1-12 carbon atoms which may be unsubstituted or substituted, e.g. by a halogen).

25

ENANTIOMERS/TAUTOMERS

In all aspects of the present invention previously discussed, the invention includes, where appropriate all enantiomers and tautomers of compounds of formula I or Ia. The man skilled in the art will recognise compounds that possess an optical properties (one or more

chiral carbon atoms) or tautomeric characteristics. The corresponding enantiomers and/or tautomers may be isolated/prepared by methods known in the art.

STEREO AND GEOMETRIC ISOMERS

5 Some of the compounds of the invention may exist as stereoisomers and/or geometric isomers – e.g. they may possess one or more asymmetric and/or geometric centres and so may exist in two or more stereoisomeric and/or geometric forms. The present invention contemplates the use of all the individual stereoisomers and geometric isomers of those
10 inhibitor agents, and mixtures thereof. The terms used in the claims encompass these forms, provided said forms retain the appropriate functional activity (though not necessarily to the same degree).

The present invention also includes all suitable isotopic variations of the agent or a
15 pharmaceutically acceptable salt thereof. An isotopic variation of an agent of the present invention or a pharmaceutically acceptable salt thereof is defined as one in which at least one atom is replaced by an atom having the same atomic number but an atomic mass different from the atomic mass usually found in nature. Examples of isotopes that can be incorporated into the agent and pharmaceutically acceptable salts thereof include isotopes
20 of hydrogen, carbon, nitrogen, oxygen, phosphorus, sulphur, fluorine and chlorine such as ^2H , ^3H , ^{13}C , ^{14}C , ^{15}N , ^{17}O , ^{18}O , ^{31}P , ^{32}P , ^{35}S , ^{18}F and ^{36}Cl , respectively. Certain isotopic variations of the agent and pharmaceutically acceptable salts thereof, for example, those in which a radioactive isotope such as ^3H or ^{14}C is incorporated, are useful in drug and/or substrate tissue distribution studies. Tritiated, i.e., ^3H , and carbon-14, i.e., ^{14}C , isotopes are
25 particularly preferred for their ease of preparation and detectability. Further, substitution with isotopes such as deuterium, i.e., ^2H , may afford certain therapeutic advantages resulting from greater metabolic stability, for example, increased *in vivo* half-life or reduced dosage requirements and hence may be preferred in some circumstances. Isotopic variations of the agent of the present invention and pharmaceutically acceptable salts

thereof of this invention can generally be prepared by conventional procedures using appropriate isotopic variations of suitable reagents.

SOLVATES

- 5 The present invention also includes the use of solvate forms of the compounds of the present invention. The terms used in the claims encompass these forms.

POLYMORPHS

- 10 The invention furthermore relates to the compounds of the present invention in their various crystalline forms, polymorphic forms and (an)hydrous forms. It is well established within the pharmaceutical industry that chemical compounds may be isolated in any of such forms by slightly varying the method of purification and or isolation from the solvents used in the synthetic preparation of such compounds.

15 PRODRUGS

- The invention further includes the compounds of the present invention in prodrug form. Such prodrugs are generally compounds of formula I wherein one or more appropriate groups have been modified such that the modification may be reversed upon administration to a human or mammalian subject. Such reversion is usually performed by
- 20 an enzyme naturally present in such subject, though it is possible for a second agent to be administered together with such a prodrug in order to perform the reversion in vivo. Examples of such modifications include ester (for example, any of those described above), wherein the reversion may be carried out by an esterase etc. Other such systems will be well known to those skilled in the art.

25

ADMINISTRATION

The pharmaceutical compositions of the present invention may be adapted for oral, rectal, vaginal, parenteral, intramuscular, intraperitoneal, intraarterial, intrathecal, intrabronchial,

subcutaneous, intradermal, intravenous, nasal, buccal or sublingual routes of administration.

For oral administration, particular use is made of compressed tablets, pills, tablets, gellules,
5 drops, and capsules. Preferably, these compositions contain from 1 to 250 mg and more preferably from 10-100 mg, of active ingredient per dose.

Other forms of administration comprise solutions or emulsions which may be injected intravenously, intraarterially, intrathecally, subcutaneously, intradermally, intraperitoneally
10 or intramuscularly, and which are prepared from sterile or sterilisable solutions. The pharmaceutical compositions of the present invention may also be in form of suppositories, pessaries, suspensions, emulsions, lotions, ointments, creams, gels, sprays, solutions or dusting powders.

15 An alternative means of transdermal administration is by use of a skin patch. For example, the active ingredient can be incorporated into a cream consisting of an aqueous emulsion of polyethylene glycols or liquid paraffin. The active ingredient can also be incorporated, at a concentration of between 1 and 10% by weight, into an ointment consisting of a white wax or white soft paraffin base together with such stabilisers and preservatives as may be
20 required.

Injectable forms may contain between 10 - 1000 mg, preferably between 10 - 250 mg, of active ingredient per dose.

25 Compositions may be formulated in unit dosage form, i.e., in the form of discrete portions containing a unit dose, or a multiple or sub-unit of a unit dose.

DOSAGE

A person of ordinary skill in the art can easily determine an appropriate dose of one of the instant compositions to administer to a subject without undue experimentation. Typically, a physician will determine the actual dosage which will be most suitable for an individual patient and it will depend on a variety of factors including the activity of the specific compound employed, the metabolic stability and length of action of that compound, the age, body weight, general health, sex, diet, mode and time of administration, rate of excretion, drug combination, the severity of the particular condition, and the individual undergoing therapy. The dosages disclosed herein are exemplary of the average case.

There can of course be individual instances where higher or lower dosage ranges are merited, and such are within the scope of this invention.

Depending upon the need, the agent may be administered at a dose of from 0.01 to 30 mg/kg body weight, such as from 0.1 to 10 mg/kg, more preferably from 0.1 to 1 mg/kg body weight.

In an exemplary embodiment, one or more doses of 10 to 150 mg/day will be administered to the patient for the treatment of malignancy.

COMBINATIONS

In a particularly preferred embodiment, the one or more compounds of the invention are administered in combination with one or more other anticancer agents, for example, existing anticancer drugs available on the market. In such cases, the compounds of the invention may be administered consecutively, simultaneously or sequentially with the one or more other anticancer agents.

Anticancer drugs in general are more effective when used in combination. In particular, combination therapy is desirable in order to avoid an overlap of major toxicities, mechanism of action and resistance mechanism(s). Furthermore, it is also desirable to

administer most drugs at their maximum tolerated doses with minimum time intervals between such doses. The major advantages of combining chemotherapeutic drugs are that it may promote additive or possible synergistic effects through biochemical interactions and also may decrease the emergence of resistance in early tumor cells which would have
5 been otherwise responsive to initial chemotherapy with a single agent. An example of the use of biochemical interactions in selecting drug combinations is demonstrated by the administration of leucovorin to increase the binding of an active intracellular metabolite of 5-fluorouracil to its target, thymidylate synthase, thus increasing its cytotoxic effects.

- 10 Numerous combinations are used in current treatments of cancer and leukemia. A more extensive review of medical practices may be found in "Oncologic Therapies" edited by E. E. Vokes and H. M. Golomb, published by Springer.

Beneficial combinations may be suggested by studying the growth inhibitory activity of
15 the test compounds with agents known or suspected of being valuable in the treatment of a particular cancer initially or cell lines derived from that cancer. This procedure can also be used to determine the order of administration of the agents, i.e. before, simultaneously, or after delivery. Such scheduling may be a feature of all the cycle acting agents identified herein.

20

NATURAL/UNNATURAL AMINO ACIDS

In one preferred embodiment of the invention, R^9 , R^{10} or R^{11} may be a natural or unnatural amino acid.

- 25 As used herein, the term "unnatural amino acid" refers to a derivative of an amino acid and may for example include alpha and alpha-disubstituted amino acids, N-alkyl amino acids, lactic acid, halide derivatives of natural amino acids such as trifluorotyrosine, p-Cl-phenylalanine, p-Br-phenylalanine, p-I-phenylalanine, L-allyl-glycine, β -alanine, L- α -amino butyric acid, L- γ -amino butyric acid, L- α -amino isobutyric acid, L- ϵ -amino caproic

acid, 7-amino heptanoic acid, L-methionine sulfone, L-norleucine, L-norvaline, p-nitro-L-phenylalanine, L-hydroxyproline, L-thioprolin, methyl derivatives of phenylalanine (Phe) such as 4-methyl-Phe, pentamethyl-Phe, L-Phe (4-amino), L-Tyr (methyl), L-Phe (4-isopropyl), L-Tic (1,2,3,4-tetrahydroisoquinoline-3-carboxyl acid), L-diaminopropionic acid and L-Phe (4-benzyl).

DEVICES

In one preferred embodiment of the invention, the R^9 , R^{10} or R^{11} groups allow for the immobilisation of the 2-phenylamino-4-heteroaryl-pyrimidine compounds onto a substrate.

10 By way of example, the R^9 , R^{10} or R^{11} groups may contain chemical functions that can be used for covalent attachment to solid phases such as functionalised polymers (*e.g.* agarose, polyacrylamide, polystyrene *etc.*) as commonly found in matrices (microtitre plate wells, microbeads, membranes, *etc.*), or used for biochemical assays or affinity chromatography. Alternatively, the R^9 , R^{10} or R^{11} groups may be linked to other small molecules (*e.g.* biotin) or

15 polypeptides (*e.g.* antigens), which can be used for non-covalent immobilisation through binding to an immobilised receptor (*e.g.* avidin or streptavidin in the case of biotin, or a specific antibodies in the case of antigens).

ASSAYS

20 Another aspect of the invention relates to the use of a compound of formula I or Ia as defined hereinabove in an assay for identifying further candidate compounds that influence the activity of one or more CDK enzymes.

Preferably, the assay is capable of identifying candidate compounds that are capable of

25 inhibiting one or more of a CDK enzyme, GSK or a PLK enzyme.

More preferably, the assay is a competitive binding assay.

Preferably, the candidate compound is generated by conventional SAR modification of a

30 compound of the invention.

As used herein, the term "conventional SAR modification" refers to standard methods known in the art for varying a given compound by way of chemical derivatisation.

Thus, in one aspect, the identified compound may act as a model (for example, a template)
5 for the development of other compounds. The compounds employed in such a test may be free in solution, affixed to a solid support, borne on a cell surface, or located intracellularly. The abolition of activity or the formation of binding complexes between the compound and the agent being tested may be measured.

10 The assay of the present invention may be a screen, whereby a number of agents are tested. In one aspect, the assay method of the present invention is a high through-put screen.

This invention also contemplates the use of competitive drug screening assays in which neutralising antibodies capable of binding a compound specifically compete with a test
15 compound for binding to a compound.

Another technique for screening provides for high throughput screening (HTS) of agents having suitable binding affinity to the substances and is based upon the method described in detail in WO 84/03564.

20 It is expected that the assay methods of the present invention will be suitable for both small and large-scale screening of test compounds as well as in quantitative assays.

Preferably, the competitive binding assay comprises contacting a compound of formula I
25 or Ia with a CDK enzyme in the presence of a known substrate of said CDK enzyme and detecting any change in the interaction between said CDK enzyme and said known substrate.

A further aspect of the invention provides a method of detecting the binding of a ligand to
30 a CDK enzyme, said method comprising the steps of:

48

- (i) contacting a ligand with a CDK enzyme in the presence of a known substrate of said CDK enzyme;
 - (ii) detecting any change in the interaction between said CDK enzyme and said known substrate;
- 5 and wherein said ligand is a compound of formula I or Ia.

One aspect of the invention relates to a process comprising the steps of:

- (a) performing an assay method described hereinabove;
- (b) identifying one or more ligands capable of binding to a ligand binding domain; and
- 10 (c) preparing a quantity of said one or more ligands.

Another aspect of the invention provides a process comprising the steps of:

- (a) performing an assay method described hereinabove;
- (b) identifying one or more ligands capable of binding to a ligand binding domain; and
- 15 (c) preparing a pharmaceutical composition comprising said one or more ligands.

Another aspect of the invention provides a process comprising the steps of:

- (a) performing an assay method described hereinabove;
- (b) identifying one or more ligands capable of binding to a ligand binding domain;
- 20 (c) modifying said one or more ligands capable of binding to a ligand binding domain;
- (d) performing the assay method described hereinabove;
- (e) optionally preparing a pharmaceutical composition comprising said one or more ligands.

- 25 The invention also relates to a ligand identified by the method described hereinabove.

Yet another aspect of the invention relates to a pharmaceutical composition comprising a ligand identified by the method described hereinabove.

Another aspect of the invention relates to the use of a ligand identified by the method described hereinabove in the preparation of a pharmaceutical composition for use in the treatment of proliferative disorders.

- 5 The above methods may be used to screen for a ligand useful as an inhibitor of one or more CDK enzymes.

The present invention is further described by way of example.

EXAMPLES**Example 1**

Chemical synthesis. The covalent attachment of solubilising moieties can be achieved in a number of different ways known in the art (Wermuth CG. Preparation of water-soluble compounds by covalent attachment of solubilizing moieties. In: Practice of Medicinal Chemistry; Academic Press: London, UK, 1996; pp 755-776). For example, amino substituents in 2-phenylamino-4-heteroaryl-pyrimidine derivatives, or their synthetic precursors, can be acylated or alkylated with carbonyl functions in appropriate solubilising moiety precursors. Similarly, carbonyl groups in the 2-phenylamino-4-heteroaryl-pyrimidine derivatives can be aminated or alkylated with appropriate solubilising moiety precursors. Halogen groups on aromatic C in phenylamino-4-heteroaryl-pyrimidines or precursors can be substituted through nucleophilic groups in solubilising moiety precursors. Suitable 2-phenylamino-4-heteroaryl-pyrimidine precursors may be prepared in accordance with the teachings of Fischer *et al* (Fischer PM, Wang S. PCT Intl. Patent Appl. Publ. WO 01/072745; Cyclacel Limited, UK, 2001). Some synthetic and analytical details for example compounds of the present invention (refer **Table 1**) are given in **Example 2** below.

Example 2

[4-(2-Amino-4-methyl-thiazol-5-yl)-pyrimidin-2-yl]-(4-morpholin-4-yl-phenyl)-amine [1]. By condensation between *N*'-[5-(3-dimethylamino-acryloyl)-4-methyl-thiazol-2-yl]-*N,N*-dimethyl-formamidine (prepared from 1-(2-amino-4-methyl-thiazol-5-yl)-ethanone and *N,N*-dimethylformamide dimethylacetal) and *N*-(4-morpholin-4-yl-phenyl)-guanidine nitrate. Yellow solid. M.p. 300 – 304 °C: ¹H-NMR (DMSO-d₆) δ: 2.46 (s, 3H, CH₃), 3.07 (m, 4H, CH₂), 3.76 (m, 4H, CH₂), 6.85 (d, 1H, *J* = 5.3 Hz, pyrimidinyl-H), 6.92 (m, 2H, Ph-H), 7.53 (br. s, 1H, NH), 7.67 (m, 2H, Ph-H), 8.30 (d, 1H, *J* = 5.4 Hz, pyrimidinyl-H), 9.25 (br. s, 1H, NH). MS (ESI⁺) *m/z* 369 [M+H]⁺ (C₁₈H₂₀N₆OS requires 368.5).

[4-(2,4-Dimethyl-thiazol-5-yl)-pyrimidin-2-yl]-(4-morpholin-4-yl-phenyl)-amine [2].

By condensation between 3-dimethylamino-1-(2,4-dimethyl-thiazol-5-yl)-propenone and *N*-(4-morpholin-4-yl-phenyl)-guanidine nitrate. Pale solid. ¹H-NMR (CDCl₃) δ: 2.69 (s, 3H, CH₃), 2.70 (s, 3H, CH₃), 3.14 (t, 4H, *J* = 4.8 Hz, CH₂), 3.72 (t, 4H, *J* = 4.9 Hz, CH₂), 5 6.89 (d, 1H, *J* = 5.1 Hz, pyrimidinyl-H), 6.95 (d, 2H, *J* = 8.8 Hz, Ph-H), 6.98 (br. s, 1H, NH), 7.53 (d, 2H, *J* = 9.1 Hz, Ph-H), 8.38 (d, 1H, *J* = 5.1 Hz, pyrimidinyl-H). MS (ESI⁺) *m/z* 368 [M+H]⁺ (C₁₉H₂₁N₅OS requires 367.5).

[4-(2-*N*-Methylamino-4-methyl-thiazol-5-yl)-pyrimidin-2-yl]-(4-morpholinophenyl)-amine

10 [3]. By condensation between 3-dimethylamino-1-(4-methyl-2-methylaminothiazol-5-yl)-propenone (prepared from 1-(4-methyl-2-methylamino-thiazol-5-yl)-ethanone and *N,N*-dimethylformamide dimethylacetal) and *N*-(4-morpholin-4-yl-phenyl)-guanidine nitrate. Pale solid. Anal. RP-HPLC: *t_R* = 10.8 min (0 – 60 % MeCN in 0.1 % aq CF₃COOH over 20 min, 1 mL/min, purity > 95 %). ¹H-NMR (DMSO-*d*₆) δ: 2.83 (s, 3H, CH₃), 2.84 (s, 3H, 15 CH₃), 3.01 (t, 4H, *J* = 5.0 Hz, CH₂), 3.72 (t, 4H, *J* = 5.0 Hz, CH₂), 6.81 (d, 2H, *J* = 5.5 Hz, pyrimidinyl-H), 6.87 (m, 2H, Ph-H), 7.61 (m, 2H, Ph-H), 8.12 (br. s, 1H, NH), 8.26 (d, 1H, *J* = 5.5 Hz, pyrimidinyl-H), 9.19 (br. s, 1H, NH). MS (ESI⁺) *m/z* 383 [M+H]⁺ (C₁₉H₂₂N₆OS requires 382.5).

20 [4-(2-Ethylamino-4-methyl-thiazol-5-yl)-pyrimidin-2-yl]-(4-morpholin-4-yl-phenyl)-amine

[4]. By condensation between 3-dimethylamino-1-(2-ethylamino-4-methyl-thiazol-5-yl)-propenone (prepared from 1-(2-ethylamino-4-methyl-thiazol-5-yl)-ethanone and *N,N*-dimethylformamide dimethylacetal) and *N*-(4-morpholin-4-yl-phenyl)-guanidine nitrate. Pale solid. Anal. RP-HPLC: *t_R* = 19.4 min (0 – 60 % MeCN in 0.1 % aq CF₃COOH over 20 25 min, 1 mL/min, purity > 95 %). ¹H-NMR (DMSO-*d*₆) δ: 1.16 (t, *J* = 7.5 Hz, 3H, CH₃), 2.48 (s, 3H, CH₃), 3.26 (m, 2H, CH₂), 3.01 (t, 4H, *J* = 5.0 Hz, CH₂), 3.72 (t, 4H, *J* = 5.0 Hz, CH₂), 6.80 (d, 2H, *J* = 5.5 Hz, pyrimidinyl-H), 6.86 (d, 2H, *J* = 9.0 Hz, Ph-H), 7.60 (d, 2H, *J* = 9.0 Hz, Ph-H), 8.25 (d, 1H, *J* = 5.0 Hz, pyrimidinyl-H), 8.50 (s, 1H, NH), 9.16 (br. s, 1H, NH).

1-(4-{4-[4-(2,4-Dimethyl-thiazol-5-yl)-pyrimidin-2-ylamino]-phenyl}-piperazin-1-yl)-ethanone [5]. A solution of 1-fluoro-4-nitrobenzene (6.7 g, 47.5 mmol), 1-piperazin-1-yl-ethanone (6.7 g, 52.3 mmol) and K₂CO₃ (6.6 g, 47.5 mmol) in DMSO (60 mL) was heated at 100 °C for 18 h. After cooling, the mixture was poured into H₂O (0.5 L). The resulting yellow precipitate was filtered and washed with H₂O to afford of 1-[4-(4-nitro-phenyl)-piperazin-1-yl]-ethanone (11.9 g). This was partially dissolved in EtOH (100 mL) and AcOH (50 mL). The mixture was warmed to *ca.* 65 °C and iron powder (-325 mesh, 12.0 g, 215 mmol) was added in 1-g portions. The mixture was heated at reflux for 2hr and filtered through a pad of Celite. The filtrate was evaporated to leave a black oil, which was basified by addition of 2 M aq NaOH and was extracted with EtOAc. The combined organics were washed with brine, dried on MgSO₄, filtered, and evaporated *in vacuo* to afford 1-[4-(4-amino-phenyl)-piperazin-1-yl]-ethanone (6.7 g) as a yellow solid. An aliquot of this material (2.0 g, 9.12 mmol) was dissolved in EtOH (5 mL) and HNO₃ was added (69 % aq soln., 1.26 mL, 19.61 mmol), followed by cyanamide (50 % w/v aq soln., 2.48 mL, 31.92 mmol). The resulting mixture was heated at reflux for 18 h. It was cooled to room temperature and poured into Et₂O (100 mL). The ethereal layer was separated and concentrated. The resulting precipitate was filtered and washed with iPrOH/Et₂O, then neat Et₂O. The light brown solid was dried to afford *N*-[4-(4-acetyl-piperazin-1-yl)-phenyl]-guanidine nitrate (1.2 g). This material (1.1g, 2.84 mmol) was dissolved in 2-methoxyethanol (14 mL) and K₂CO₃ (0.79 g, 5.68 mmol) was added, followed by 3-dimethylamino-1-(2,4-dimethyl-thiazol-5-yl)-propenone (0.60 g, 2.84 mmol). The resulting mixture was heated at 115 °C for 18 h. It was cooled and concentrated. The residue was purified by SiO₂ chromatography (9:1 EtOAc / 2 M NH₃ in MeOH) and recrystallisation from iPr₂O/MeOH to afford the title compound (930 mg) as a light brown solid. ¹H-NMR (CDCl₃) δ: 2.15 (s, 3H, CH₃), 2.69 (s, 3H, CH₃), 2.71 (s, 3H, CH₃), 3.12 (t, 2H, *J* = 5.4 Hz, CH₂), 3.15 (t, 2H, *J* = 5.4 Hz, CH₂), 3.63 (t, 2H, *J* = 5.4 Hz, CH₂), 3.79 (t, 2H, *J* = 5.4 Hz, CH₂), 6.90 (d, 1H, *J* = 5.4 Hz, pyrimidinyl-H), 6.96 (m, 2H, Ph-H), 6.98 (br. s, 1H, NH), 7.54 (m, 2H, Ph-H), 8.39 (d, 1H, *J* = 5.4 Hz, pyrimidinyl-H). MS (ESI⁺) *m/z* 409.6 (C₂₁H₂₄N₆OS requires 408.5).

[4-(2,4-Dimethyl-thiazol-5-yl)-pyrimidin-2-yl]-(4-piperazin-1-yl-phenyl)-amine [6]. To a solution of 1-(4-{4-[4-(2,4-dimethyl-thiazol-5-yl)-pyrimidin-2-ylamino]-phenyl}-piperazin-1-yl)-ethanone (0.67 g, 1.64 mmol) in EtOH (3 mL) was added 2 M aq HCl (25 mL) in a steady stream. The mixture was heated at reflux for 1 h, cooled, and basified by addition of solid Na₂CO₃. The product was extracted with EtOAc. The combined organics were washed with brine, dried on Na₂SO₄, filtered, and evaporated to afford the title compound (580 mg) as a yellow solid. ¹H-NMR (CDCl₃) τ : 2.62 (s, 3H, CH₃), 2.63 (s, 3H, CH₃), 2.99 (m, 4H, CH₂), 3.06 (m, 4H, CH₂), 6.81 (d, 1H, *J* = 5.4 Hz, pyrimidinyl-H), 6.89 (m, 3H, Ph-H, NH), 7.44 (m, 2H, Ph-H), 8.31 (d, 1H, *J* = 5.4 Hz, pyrimidinyl-H). MS (ESI⁺) *m/z* 367 (C₁₉H₂₂N₆S requires 366.5).

[4-(2,4-Dimethyl-thiazol-5-yl)-pyrimidin-2-yl]-[4-(4'-2''-ethoxyethanolpiperazino)-phenyl]-amine [7]. A mixture of [4-(2,4-dimethyl-thiazol-5-yl)-pyrimidin-2-yl]-(4-piperazin-1-yl-phenyl)-amine (0.1 g, 0.27 mmol), 2-(2-chloro-ethoxy)-ethanol (35 μ L, 0.33 mmol), NaI (49 mg, 0.33 mmol), and K₂CO₃ (37 mg, 0.27 mmol) in MeCN (2 mL) in a sealed tube was heated at 170 °C for 15 min in a microwave reactor (SmithCreator, Personal Chemistry Ltd). The solvent was evaporated to dryness and the residue was purified by SiO₂ chromatography (98:2 to 95:5 EtOAc / 2 M NH₃ in MeOH) to afford the title compound (78 mg) as yellow foam. ¹H-NMR (CDCl₃) δ : 2.69 (s, 3H, CH₃), 2.71 (s, 3H, CH₃), 2.80-2.91 (m, 6H, CH₂), 3.30 (m, 4H, CH₂), 3.67 (m, 2H, CH₂), 3.73 (m, 2H, CH₂), 3.79 (m, 2H, CH₂), 6.89 (d, 1H, *J* = 5.4 Hz, pyrimidinyl-H), 6.97 (m, 3H, Ph-H & NH), 7.52 (m, 2H, Ph-H), 8.02 (br. s, 1H, OH), 8.38 (d, 1H, *J* = 5.4 Hz, pyrimidinyl-H). MS (ESI⁺) *m/z* 456 (C₂₃H₃₀N₆O₂S requires 454.6).

3-(4-{4-[4-(2,4-Dimethyl-thiazol-5-yl)-pyrimidin-2-ylamino]-phenyl}-piperazin-1-yl)-propan-1-ol [8]. Yellow solid. ¹H-NMR (CDCl₃) δ : 2.69 (s, 3H, CH₃), 2.71 (s, 3H, CH₃), 2.75 (m, 2H, CH₂), 3.21 (m, 2H, CH₂), 3.84 (t, 2H, *J* = 5.1 Hz, CH₂), 6.89 (d, 1H, *J* = 5.4 Hz, pyrimidinyl-H), 6.95 (m, 2H, Ph-H), 6.98 (br. s, 1H, NH), 7.51 (m, 2H, Ph-H), 8.38 (d, 1H, *J* = 5.4 Hz, pyrim-H). MS (ESI⁺): *m/z* 425.8 (C₂₂H₂₈N₆OS requires 424.6).

2-(4-{4-[4-(2,4-Dimethyl-thiazol-5-yl)-pyrimidin-2-ylamino]-phenyl}-piperazin-1-yl)-ethanol [9]. Yellow solid. ¹H-NMR (CDCl₃) δ: 2.55 (t, 2H, *J* = 5.4 Hz, CH₂), 2.62 (m, 10H, CH₃ & CH₂), 3.12 (t, 4H, *J* = 4.9 Hz, CH₂), 3.60 (t, 2H, *J* = 5.4 Hz, CH₂), 6.81 (d, 1H, *J* = 5.4 Hz, pyrimidinyl-H), 6.88 (m, 2H, Ph-H), 7.05 (br. s, 1H, NH), 7.45 (m, 2H, Ph-H), 8.30 (d, 1H, *J* = 5.1 Hz, pyrimidinyl-H). MS (ESI⁺) *m/z* 411.7 (C₂₁H₂₆N₆OS requires 410.5).

[4-(2,4-Dimethyl-thiazol-5-yl)-pyrimidin-2-yl]-[4-(4-methanesulfonyl-piperazin-1-yl)-phenyl]-amine [10]. A mixture of [4-(2,4-dimethyl-thiazol-5-yl)-pyrimidin-2-yl]-(4-piperazin-1-yl-phenyl)-amine (86 mg, 0.23 mmol) in anhydrous CH₂Cl₂ (2 mL) was added Et₃N (39 μL, 0.28 mmol). After cooling to 0 °C, methanesulfonyl chloride (22 μL, 0.28 mmol) was added dropwise. After 15 min stirring, the reaction mixture was warmed to room temperature and stirring was continued for 18 h. After evaporation, the residue was purified by SiO₂ chromatography (98:2 to 95:5 EtOAc / 2 M NH₃ in MeOH) to afford the title compound (61 mg) as a yellow solid. ¹H-NMR (CDCl₃) δ: 2.69 (s, 3H, CH₃), 2.71 (s, 3H, CH₃), 2.84 (s, 3H, CH₃), 3.26 (t, 4H, *J* = 5.1 Hz, CH₂), 3.41 (t, 4H, *J* = 5.1 Hz, CH₂), 6.91 (d, 1H, *J* = 5.1 Hz, pyrimidinyl-H), 6.98 (d, 2H, *J* = 8.8 Hz, Ph-H), 7.10 (br. s, 1H, NH), 7.56 (d, 2H, *J* = 8.8 Hz, Ph-H), 8.30 (d, 1H, *J* = 5.1 Hz, pyrimidinyl-H). MS (ESI⁺) *m/z* 446 (C₂₀H₂₄N₆O₂S₂ requires 444.6).

20

[4-(4-Benzyl-piperazin-1-yl)-phenyl]-[4-(4-methyl-2-methylamino-thiazol-5-yl)-pyrimidin-2-yl]-amine [11]. 4-(4-Benzyl-piperazin-1-yl)-phenylamine (2.17 g, 8.12 mmol) was partially dissolved in EtOH (5 mL) and HNO₃ (69 % aq soln., 1.05 mL, 16.32 mmol) was added dropwise, followed by cyanamide (50 % aq soln., 1.13 mL, 16.32 mmol). The mixture was heated for 18 h at reflux. After working up *N*-[4-(4-benzyl-piperazin-1-yl)-phenyl]-guanidine nitrate (1.16 g) was obtained as a purple solid. A mixture of this material (2.66 mmol), 3-dimethylamino-1-(4-methyl-2-methylaminothiazol-5-yl)-propanone (0.60 g, 2.66 mmol), and K₂CO₃ (0.74 g, 5.32 mmol) in 2-methoxyethanol (15 mL) was heated at 120 °C for 18 h. After cooling, it was poured into EtOAc (100 mL) and filtered through a pad of silica. The filtrate was evaporated and the residue was purified by

30

SiO₂ chromatography (heptane/EtOAc) to afford the title compound (442 mg) as a light tan solid. ¹H-NMR (CD₃OD) δ: 2.44 (s, 3H, CH₃), 2.56-2.58 (m, 4H, CH₂), 2.91 (s, 3H, CH₃), 3.09 (m, 4H, CH₂), 3.51 (s, 2H, CH₂), 6.70 (d, 1H, *J* = 5.6 Hz, pyrimidinyl-H), 6.87 (m, 2H, Ph-H), 7.22 (m, 1H, Ph-H), 7.27 (m, 4H, Ph-H), 7.43 (m, 2H, Ph-H), 8.15 (d, 1H, *J* = 5.4 Hz, pyrimidinyl-H). MS (ESI⁺) *m/z* 473.2 (C₂₆H₂₉N₇S requires 471.6).

[4-(2,4-Dimethyl-thiazol-5-yl)-pyrimidin-2-yl]-[4-(4-methyl-piperazin-1-yl)-phenyl]-amine [12]. By condensation between 3-dimethylamino-1-(2,4-dimethyl-thiazol-5-yl)-propenone and *N*-[4-(4-methyl-piperazin-1-yl)-phenyl]-guanidine nitrate. Light yellow solid. ¹H-NMR (CDCl₃) δ: 2.37 (s, 3H, CH₃), 2.61 (m, 4H, CH₂), 2.69 (s, 3H, CH₃), 2.70 (s, 3H, CH₃), 3.20 (m, 4H, CH₂), 6.88 (d, 1H, *J* = 5.1 Hz, pyrimidinyl-H), 6.94 (s, 1H, NH), 6.96 (d, 2H, *J* = 8.8 Hz, Ph-H), 7.51 (d, 2H, *J* = 8.8 Hz, Ph-H), 8.38 (d, 1H, *J* = 5.1 Hz, pyrimidinyl-H).

3-Amino-*N*-[4-methyl-5-[2-(3-nitro-phenylamino)-pyrimidin-4-yl]-thiazol-2-yl]-propionamide [13]. A mixture of [4-(2-amino-4-methyl-thiazol-5-yl)-pyrimidin-2-yl]-(3-nitro-phenyl)-amine (0.12 g, 0.37 mmol), Boc-βAla-OH (0.18 g, 0.93 mmol), 1,3-diisopropylcarbodiimide (0.07 mL, 0.45 mmol) and 4-*N,N*-dimethylaminopyridine (36 mg, 0.3 mmol) in of dry DMF (2 mL) was stirred at room temperature for 24 h. The reaction mixture was poured into ice water and extracted with EtOAc. The organics were combined and washed with brine, filtered, and dried on MgSO₄. The solvent was evaporated to give brown residue, which was purified by SiO₂ chromatography (1:1 EtOAc/PE) to afford ({4-Methyl-5-[2-(3-nitro-phenylamino)-pyrimidin-4-yl]-thiazol-2-ylcarbamoyl}-methyl)-carbamic acid tert-butyl ester as light yellow solid. Anal. RP-HPLC: *t_R* = 19.5 min (0 – 60 % MeCN in 0.1 % aq CF₃COOH over 20 min, 1 mL/min, purity >93 %). ¹H-NMR (DMSO-*d*₆) δ: 1.11 (s, 9H, CH₃), 2.54 (s, 3H, CH₃), 3.62 (m, 2H, CH₂), 5.38 (m, 2H, CH₂), 7.06 (d, 1H, *J* = 5.5 Hz, pyrimidinyl-H), 7.46 (m, 1H, Ph-H), 7.72 (m, 1H, Ph-H), 8.15 (m, 1H, Ph-H), 8.45 (d, 1H, *J* = 5.2 Hz, pyrimidinyl-H), 8.77 (s, 1H, NH). A solution of this material (97 mg, 0.19 mmol) in dioxane (5 mL) was treated with CF₃COOH (1 mL). After stirring at room temperature for 22 h, the reaction mixture was evaporated and purified by

preparative RP-HPLC (0 – 60 % MeCN in 0.1 % aq CF₃COOH over 40 min, 9 mL/min) to afford the titled compound (34 mg) as a light yellow solid. Anal. RP-HPLC: *t_R* = 14.0 min (0 – 60 % MeCN in 0.1 % aq CF₃COOH over 20 min, 1 mL/min, purity >93 %). ¹H-NMR (CD₃OD) δ: 2.64 (s, 3H, CH₃), 3.30 (m, 2H, CH₂), 3.33 (m, 2H, CH₂), 7.14 (d, 1H, *J* = 5.5 Hz, pyrimidinyl-H), 7.52 (t, 1H, *J* = 8.2 Hz, Ph-H), 7.63 (m, 1H, Ph-H), 7.96 (m, 1H, Ph-H), 8.46 (d, 1H, *J* = 5.5 Hz, pyrimidinyl-H), 8.88 (s, 1H, NH). MS (ESI⁺) *m/z* 400.6 (C₁₇H₁₇N₇O₃S requires 399.4).

(2*S*)-2-Amino-3-hydroxy-N-{4-methyl-5-[2-(3-nitro-phenylamino)-pyrimidin-4-yl]-thiazol-2-yl}-propionamide [14]. From Boc-L-Ser-OH. Pale solid. Anal. RP-HPLC: *t_R* = 13.7 min (0 – 60 % MeCN in 0.1 % aq CF₃COOH over 20 min, 1 mL/min, purity > 93 %). ¹H-NMR (CD₃OD) δ: 2.45 (s, 3H, CH₃), 3.65-3.77 (m, 3H, CH & CH₂), 3.95 (br. s, 1H, OH), 7.03 (d, 1H, *J* = 5.5 Hz, pyrimidinyl-H), 7.36 (t, 1H, *J* = 6.5 Hz, Ph-H), 7.63 (m, 1H, Ph-H), 7.97 (m, 1H, Ph-H), 8.41 (d, 1H, *J* = 5.5 Hz, pyrimidinyl-H), 8.63 (s, 1H, NH).

(2*R*,3*R*)-2-Amino-3-hydroxy-N-{4-methyl-5-[2-(3-nitro-phenylamino)-pyrimidin-4-yl]-thiazol-2-yl}-butyramide [15]. From Boc-D-Thr-OH. Yellow solid. ¹H-NMR (CD₃OD) δ: 1.35 (d, 3H, *J* = 6.1 Hz, CH₃), 2.66 (s, 1H, CH), 3.95 (d, 1H, *J* = 5.4 Hz, CH), 7.17 (d, 1H, *J* = 5.4 Hz, pyrimidinyl-H), 7.51 (m, 1H, Ph-H), 7.97 (m, 1H, Ph-H), 8.51 (d, 1H, *J* = 5.5 Hz, pyrimidinyl-H), 8.92 (m, 1H, Ph-H).

(2*R*)-2-Amino-N-{4-methyl-5-[2-(3-nitro-phenylamino)-pyrimidin-4-yl]-thiazol-2-yl}-butyramide [16]. From Boc-D-Abu-OH. Yellow solid. ¹H-NMR (DMSO-*d*₆) δ: 0.93 (t, 3H, *J* = 7.6 Hz, CH₃), 0.94 (m, 2H, CH₂), 1.60 (s, 3H, CH₃), 3.00 (t, 1H, *J* = 6.9 Hz, CH), 6.10 (d, 1H, *J* = 5.5 Hz, pyrimidinyl-H), 6.44 (m, 1H, Ph-H), 6.77 (m, 1H, Ph-H), 6.87 (m, 1H, Ph-H), 7.42 (d, 1H, *J* = 5.5 Hz, pyrimidinyl-H) and 7.86 (br. s, 1H, NH). MS (ESI⁺) *m/z* 414.6 (C₁₈H₁₉N₇O₃S requires 413.5).

(2*S*,3*S*)-2-Amino-3-hydroxy-N-{4-methyl-5-[2-(3-nitro-phenylamino)-pyrimidin-4-yl]-thiazol-2-yl}-butyramide [17]. From Boc-L-Thr-OH. Yellow solid. ¹H-NMR (CD₃OD) δ:

57

1.19 (d, 3H, $J = 6.6$ Hz, CH₃), 2.63 (s, 3H, CH₃), 3.89 (m, 1H, CH), 4.10 (m, 1H, CH), 7.20 (d, 1H, $J = 5.5$ Hz, pyrimidinyl-H), 7.55 (t, 1H, $J = 8.0$ Hz, Ph-H), 7.81 (d, 1H, $J = 8.5$ Hz, Ph-H), 8.16 (d, 1H, $J = 8.5$ Hz, Ph-H), 8.58 (d, 1H, $J = 5.5$ Hz, pyrimidinyl-H), 8.80 (s, 1H, Ph-H) and 10.17 (s, 1H, NH).

5

4-Amino-N-{4-methyl-5-[2-(3-nitro-phenylamino)-pyrimidin-4-yl]-thiazol-2-yl}-

butyramide [18]. Yellow solid. ¹H-NMR (CD₃OD) δ : 1.87 (m, 2H, CH₂), 2.52 (s, 3H, CH₃), 2.56 (m, 2H, CH₂), 2.84 (m, 2H, CH₂), 7.17 (d, 1H, $J = 5.3$ Hz, pyrimidinyl-H), 7.55 (t, 1H, $J = 8.3$ Hz, Ph-H), 7.79 (m, 1H, Ph-H), 8.13 (m, 1H, Ph-H), 8.54 (d, 1H, $J = 5.3$ Hz, pyrimidinyl-H), 8.83 (s, 1H, Ph-H) and 10.12 (s, 1H, NH).

10

3-Amino-N-{4-methyl-5-[2-(4-morpholin-4-yl-phenylamino)-pyrimidin-4-yl]-thiazol-2-yl}-

propionamide [19]. Yellow solid. ¹H-NMR (DMSO-d₆) δ : 2.57 (m, 2H, CH₂), 2.71 (s, 3H, CH₃), 2.92 (m, 2H, CH₂), 3.02 (m, 4H, CH₂), 3.73 (m, 4H, CH₂), 6.87 (d, 2H, $J = 8.0$ Hz, Ph-H), 6.95 (d, 1H, $J = 5.3$ Hz, pyrimidinyl-H), 7.61 (d, 1H, $J = 8.0$ Hz, Ph-H), 8.38 (d, 1H, $J = 5.5$ Hz, pyrimidinyl-H) and 9.32 (s, 1H, NH). MS (ESI⁺) m/z 439 (C₂₁H₂₅N₇O₂S requires 439.5).

15

3-Bromo-N-{5-[2-(4-fluoro-phenylamino)-pyrimidin-4-yl]-4-methyl-thiazol-2-yl}-

propionamide [20]. A solution of [4-(2-amino-4-methyl-thiazol-5-yl)-pyrimidin-2-yl]-(4-fluoro-phenyl)-amine (0.3 g, 1.0 mmol) in DMF (2 mL) was cooled on an ice bath and was treated with 2-bromopropionyl chloride (0.17 g, 1.0 mmol). After completion of the addition, the reaction mixture was allowed to stir at room temperature for 18 h. It was poured into ice water and was extracted with CH₂Cl₂. The organics were combined, washed with brine, dried on MgSO₄, and the solvent was evaporated to leave a brown residue. This was purified by SiO₂ chromatography (1:1 EtOAc/PE) to afford the title compound as a light yellow solid. Anal. RP-HPLC: $t_R = 17.1$ min (0 – 60 % MeCN in 0.1 % aq CF₃COOH over 20 min, 1 mL/min, purity > 93 %). ¹H-NMR (DMSO-d₆) δ : 1.99 (m, 2H, CH₂), 2.68 (s, 3H, CH₃), 4.65 (m, 2H, CH₂), 7.02 (d, 1H, $J = 5.2$ Hz, pyrimidinyl-H), 7.11 (m, 2H, Ph-H), 7.62 (m, 2H, Ph-H), 8.19 (d, 1H, $J = 5.5$ Hz, pyrimidinyl-H).

30

N-{5-[2-(4-Fluoro-phenylamino)-pyrimidin-4-yl]-4-methyl-thiazol-2-yl}-3-morpholin-4-yl-propionamide [21]. A solution of 3-bromo-N-{5-[2-(4-fluoro-phenylamino)-pyrimidin-4-yl]-4-methyl-thiazol-2-yl}-propionamide (20 mg, 0.046 mmol) and morpholine (8 μ L, 0.092 mmol) in DMF (2 mL) was stirred at room temperature for 2 h. The reaction mixture was purified by preparative RP-HPLC (0 – 60 % MeCN in 0.1 % aq CF₃COOH over 40 min, 9 mL/min) to afford the title compound as a pale solid. Anal. RP-HPLC: t_R = 13.3 min (0 – 60 % MeCN in 0.1 % aq CF₃COOH over 20 min, 1 mL/min, purity > 93 %). ¹H-NMR (CD₃OD) δ : 2.66 (s, 3H, CH₃), 3.23 (m, 2H, CH₂), 3.33 (m, 2H, CH₂), 3.41 (m, 4H, CH₂), 3.87 (m, 2H, CH₂), 4.13 (m, 2H, CH₂), 7.04-7.09 (m, 3H, pyrimidinyl-H & Ph-H), 7.68 (m, 2H, Ph-H), 8.39 (d, 1H, J = 5.5 Hz, pyrimidinyl-H). MS (ESI⁺) m/z 443.3 (C₂₁H₂₃FN₆O₂S requires 442.5).

N-{4-Methyl-5-[2-(3-nitro-phenylamino)-pyrimidin-4-yl]-thiazol-2-yl}-3-morpholin-4-yl-propionamide [22]. Yellow solid. ¹H-NMR (CD₃OD) δ : 2.64 (s, 3H, CH₃), 3.23 (m, 2H, CH₂), 3.33 (m, 2H, CH₂), 3.41 (m, 4H, CH₂), 3.87 (m, 2H, CH₂), 4.13 (m, 2H, CH₂), 7.04-7.09 (m, 3H, pyrimidinyl-H & Ph-H), 7.68 (m, 2H, Ph-H), 8.39 (d, 1H, J = 5.5 Hz, pyrimidinyl-H). MS (ESI⁺) m/z 443.3 (C₂₁H₂₃FN₆O₂S requires 442.5).

N-{4-Methyl-5-[2-(3-nitro-phenylamino)-pyrimidin-4-yl]-thiazol-2-yl}-3-(4-methyl-piperazin-1-yl)-propionamide [23]. Yellow solid. ¹H-NMR (CD₃OD) δ : 2.94 (s, 3H, CH₃), 3.01 (m, 2H, CH₂), 3.24-3.43 (m, 4H, CH₂), 3.65 (m, 2H, CH₂), 7.14 (d, 1H, J = 5.5 Hz, pyrimidinyl-H), 7.52 (m, 1H, Ph-H), 7.86 (d, 1H, J = 8.0 Hz, Ph-H), 7.99 (d, 1H, J = 8.0 Hz, Ph-H), 8.44 (m, 1H, Ph-H) and 8.79 (d, 1H, J = 5.5 Hz, pyrimidinyl-H). MS (ESI⁺) m/z 485 (C₂₂H₂₆N₈O₃S requires 482.6).

2-Chloro-N-{4-methyl-5-[2-(3-nitro-phenylamino)-pyrimidin-4-yl]-thiazol-2-yl}-acetamide [24]. A solution of [4-(2-amino-4-methyl-thiazol-5-yl)-pyrimidin-2-yl]-(3-nitro-phenyl)-amine (0.33 g, 1.0 mmol) in dry DMF (3 mL) was cooled on an ice-water bath. Chloroacetyl chloride (0.22 g, 2.0 mmol) and pyridine (80 μ L) were added. After stirring at room temperature for 18 h, the reaction mixture was concentrated, poured into ice water

and extracted with CH₂Cl₂. The organic phases were combined, washed with brine, dried on MgSO₄ and evaporated to dryness. The resulting greenish residue was purified by SiO₂ chromatography (1:1 EtOAc/PE) to afford the title compound as a gray solid. Anal. RP-HPLC: t_R = 20.6 min (0 – 60 % MeCN in 0.1 % aq CF₃COOH over 20 min, 1 mL/min, purity > 97 %). ¹H-NMR (DMSO-d₆) δ: 2.45 (s, 3H, CH₃), 4.12 (s, 2H, CH₂), 7.03 (d, 1H, J = 5.2 Hz, pyrimidinyl-H), 7.42 (m, 1H, Ph-H), 7.63 (m, 1H, Ph-H), 8.01 (m, 1H, Ph-H), 8.41 (d, 1H, J = 5.2 Hz, pyrimidinyl-H), 8.64 (s, 1H, Ph-H).

2-Chloro-N-{5-[2-(3-chloro-phenylamino)-pyrimidin-4-yl]-4-methyl-thiazol-2-yl}-acetamide [25]. Brown solid. ¹H-NMR (DMSO-d₆) δ: 2.65 (s, 3H, CH₃), 4.42 (s, 2H, CH₂), 7.01 (m, 1H, Ph-H), 7.25 (m, 1H, Ph-H), 7.61 (d, 1H, J = 5.5 Hz, pyrimidinyl-H), 7.98 (s, 1H, Ph-H), 8.75 (d, 1H, J = 5.5 Hz, pyrimidinyl-H) and 10.09 (br. s, 1H, NH).

2-Chloro-N-{5-[2-(3-methoxy-phenylamino)-pyrimidin-4-yl]-4-methyl-thiazol-2-yl}-acetamide [26]. Brown solid. ¹H NMR (DMSO-d₆) δ: 2.58 (s, 3H, CH₃), 3.78 (s, 3H, CH₃), 4.36 (s, 2H, CH₂), 6.51 (m, 1H, Ph-H), 7.08 (d, 1H, J = 5.5, pyrimidinyl-H), 7.14 (t, 1H, J = 8.0 Hz, Ph-H), 7.23 (m, 1H, Ph-H), 7.59 (s, 1H, Ph-H) and 8.45 (d, 1H, J = 5.5 Hz, pyrimidinyl-H).

2-Chloro-N-{5-[2-(4-dimethylamino-phenylamino)-pyrimidin-4-yl]-4-methyl-thiazol-2-yl}-acetamide [27]. Yellow solid. ¹H-NMR (CD₃OD) δ: 2.57 (s, 3H, CH₃), 3.23 (s, 6H, CH₃), 4.22 (s, 2H, CH₂), 7.06 (d, 1H, J = 5.5 Hz, pyrimidinyl-H), 7.57 (d, 2H, J = 9.5 Hz, Ph-H), 7.73 (d, 2H, J = 9.5 Hz, Ph-H), 8.14 (br. s, 1H, NH) and 8.35 (d, 1H, J = 5.5 Hz, pyrimidinyl-H).

4-({4-Methyl-5-[2-(3-nitro-phenylamino)-pyrimidin-4-yl]-thiazol-2-ylcarbamoyl}-methyl)-piperazine-1-carboxylic acid tert-butyl ester [28]. A solution of 2-chloro-N-{4-methyl-5-[2-(3-nitro-phenylamino)-pyrimidin-4-yl]-thiazol-2-yl}-acetamide (40 mg, 0.1 mmol) in DMF (2 mL) was cooled on an ice bath. Piperazine-1-carboxylic acid tert-butyl ester (40 mg, 0.21 mmol) was added. After stirring at room temperature for 16 h, the reaction

mixture was purified by preparative RP-HPLC (0 – 60 % MeCN in 0.1 % aq CF₃COOH over 40 min, 9 mL/min) to afford the title compound (20 mg) as a light yellow solid. Anal. RP-HPLC: t_R = 16.9 min (0 – 60 % MeCN in 0.1 % aq CF₃COOH over 20 min, 1 mL/min, purity > 97 %). ¹H-NMR (CD₃OD) δ : 2.65 (s, 3H, CH₃), 2.87 (m, 4H, CH₂), 3.36 (m, 4H, CH₂), 4.66 (s, 2H, CH₂), 7.71 (m, 1H, pyrimidinyl-H), 7.89 (t, 1H, J = 8.1 Hz, Ph-H), 8.15 (m, 1H, Ph-H), 7.42 (m, 1H, Ph-H), 7.67 (m, 2H, pyrimidinyl-H & Ph-H).

N-{5-[2-(4-Dimethylamino-phenylamino)-pyrimidin-4-yl]-4-methyl-thiazol-2-yl}-2-[2-(2-hydroxy-ethoxy)-ethylamino]-acetamide [29]. Anal. RP-HPLC: t_R = 10.52 min (0 – 60 % MeCN in 0.1 % aq CF₃COOH over 20 min, 1 mL/min, purity > 97 %). ¹H-NMR (CD₃OD) δ : 3.28 (s, 3H, CH₃), 3.41 (s, 6H, CH₃), 4.04 (m, 2H, CH₂), 4.28 (m, 2H, 4H, CH₂), 4.34 (m, 2H, CH₂), 4.50 (m, 2H, CH₂), 4.94 (br. s, 2H, CH₂), 7.92 (d, 1H, J = 5.5 Hz, pyrimidinyl-H), 8.00 (d, 2H, J = 9.0 Hz, Ph-H), 8.41 (m, 1H, J = 9.0 Hz, Ph-H), 8.87 (br. s, 1H, NH/OH), 9.31 (d, 1H, J = 5.5 Hz, pyrimidinyl-H).

6-[5-(2-Oxo-hexahydro-thieno[3,4-d]imidazol-4-yl)-pentanoylamino]-hexanoic acid (2-{4-methyl-5-[2-(4-morpholin-4-yl-phenylamino)-pyrimidin-4-yl]-thiazol-2-yl}carbamoylethyl)-amide [30]. A solution of 3-amino-*N*-{4-methyl-5-[2-(4-morpholin-4-yl-phenylamino)-pyrimidin-4-yl]-thiazol-2-yl}-propionamide (75mg, 0.17 mmol) in DMF (1 mL) was treated with succinimidyl-6-(biotinamido)hexanoate (32 mg, 0.085 mmol). After stirring at room temperature for 3 h, the reaction mixture was purified by preparative RP-HPLC (0 – 60 % MeCN in 0.1 % aq CF₃COOH over 40 min, 9 mL/min) to afford the title compound as an orange solid. Anal. RP-HPLC: t_R = 13.2 min (0 – 60 % MeCN in 0.1 % aq CF₃COOH over 20 min, 1 mL/min, purity > 97 %). MS (ESI+) m/z 776 (C₃₇H₅₀N₁₀O₅S₂ requires 778.9).

N-{5-[2-(4-Fluoro-phenylamino)-pyrimidin-4-yl]-4-methyl-thiazol-2-yl}-methanesulfonamide [31]. This compound was prepared from [4-(2-amino-4-methyl-thiazol-5-yl)-pyrimidin-2-yl]-(4-fluoro-phenyl)-amine by treatment with methylsulfonyl chloride and Et₃N in DMF. Gray solid; ¹H-NMR (DMSO-*d*₆) δ : 3.31 (s, 3H, CH₃), 3.63 (s,

61

3H, CH₃), 7.32 (m, 1H, pyrimidinyl-H), 7.42 (m, 2H, Ph-H), 8.11 (m, 2H, Ph-H), 8.63 (d, 1H, *J* = 5.0 Hz, pyrimidinyl-H).

N-{4-Methyl-5-[2-(3-nitro-phenylamino)-pyrimidin-4-yl]-thiazol-2-yl}-methane

5 *sulfonamide* [32]. A mixture of [4-(2-amino-4-methyl-thiazol-5-yl)-pyrimidin-2-yl]-(3-nitro-phenyl)-amine (1.0 mmol, 0.33 g) and methylsulfonyl chloride (2.0 mmol, 0.22 g) in dry DMF (2 mL) was added Et₃N (0.28 mL). The reaction mixture was stirred at room temperature for 20 h. After cooling, the mixture was diluted with EtOAc, washed with
10 brine, and dried over MgSO₄. The solvent was evaporated and the residue was purified by preparative RP-HPLC using a gradient from 10 – 70 % MeCN in 0.1 % aq CF₃COOH over 40 min. The title compound was obtained as an orange solid. Anal. RP-HPLC: *t_R* = 17.4 min (0 – 60 % MeCN in 0.1 % aq CF₃COOH over 20 min, 1 mL/min, purity > 97 %). ¹H-NMR (DMSO-*d*₆) δ : 3.10 (s, 3H, CH₃), 3.25 (s, 3H, CH₃), 7.05 (d, 1H, *J* = 5.2 Hz, pyrimidinyl-H), 7.42 (m, 1H, Ph-H), 7.63 (m, 1H, Ph-H), 7.98 (m, 1H, Ph-H), 8.21 (d, 1H,
15 *J* = 5.2 Hz, pyrimidinyl-H), 8.42 (s, 1H, Ph-H), 9.18 (s, 1H, NH).

2-Chloro-N-{5-[2-(4-fluoro-phenylamino)-pyrimidin-4-yl]-4-methyl-thiazol-2-yl}-

acetamide [33]. This compound was prepared from [4-(2-amino-4-methyl-thiazol-5-yl)-pyrimidin-2-yl]-(4-fluoro-phenyl)-amine. ¹H-NMR (DMSO-*d*₆) δ : 2.94 (s, 3H, CH₃), 4.75
20 (s, 2H, CH₂), 7.44 (m, 3H, pyrimidinyl-H & Ph-H), 8.09 (m, 2H, Ph-H), 8.28 (s, 1H, NH), 8.80 (d, 1H, *J* = 5.2 Hz, pyrimidinyl-H).

2-Chloro-N-{5-[2-(4-chloro-phenylamino)-pyrimidin-4-yl]-4-methyl-thiazol-2-yl}-

acetamide [34]. This compound was prepared by chloroacetylation of [4-(2-amino-4-methyl-thiazol-5-yl)-pyrimidin-2-yl]-(4-chloro-phenyl)-amine. Brown solid. ¹H-NMR (DMSO-*d*₆) δ : 2.65 (s, 3H, CH₃), 4.42 (s, 2H, CH₂), 7.01 (m, 1H, Ph-H), 7.25 (m, 1H, Ph-H), 7.61 (d, 1H, *J* = 5.5, pyrimidinyl-H), 7.98 (s, 1H, Ph-H), 8.75 (d, 1H, *J* = 5.5 Hz, pyrimidinyl-H), 10.09 (br. s, 1H, NH).

N-{5-[2-(4-Chloro-phenylamino)-pyrimidin-4-yl]-4-methyl-thiazol-2-yl}-3-morpholin-4-yl-propionamide [35]. By treatment of 3-chloro-*N*-{5-[2-(4-chloro-phenylamino)-pyrimidin-4-yl]-4-methyl-thiazol-2-yl}-propionamide with morpholine. Anal. RP-HPLC: t_R = 12.7 min (10 – 70 % MeCN, purity > 95 %). $^1\text{H-NMR}$ (CDCl_3) δ : 1.50 (m, 2H, CH_2), 2.52 (s, 3H, CH_3), 3.05-3.78 (m, 8H, CH_2), 3.81 (m, 2H, CH_2), 7.12 (d, 1H, J = 5.5Hz, pyrimidinyl-H), 7.30 (d, 2H, J = 7.0Hz, Ph-H), 7.80 (d, 2H, J = 7.0Hz, Ph-H), 8.51 (d, 1H, J = 5.0Hz, pyrimidinyl-H), 9.80 (brs, 1H, NH).

N-{5-[2-(4-Chloro-phenylamino)-pyrimidin-4-yl]-4-methyl-thiazol-2-yl}-3-(2-diethylamino-ethylamino)-propionamide [36]. By treatment of 3-bromo-*N*-{5-[2-(4-chloro-phenylamino)-pyrimidin-4-yl]-4-methyl-thiazol-2-yl}-propionamide with N^1, N^1 -diethyl-ethane-1,2-diamine. Anal. RP-HPLC: t_R = 11.8 min (10 – 70 % MeCN, purity > 97 %). $^1\text{H-NMR}$ (DMSO-D_6) δ : 1.20 (t, 6H, J = 7.0Hz, CH_3), 1.53 (d, 2H, J = 6.5Hz, CH_2), 2.52 (s, 6H, CH_3), 3.18 (m, 4H, CH_2), 3.28 (m, 4H, CH_2), 7.13 (d, 1H, J = 5.0Hz, pyrimidinyl-H), 7.30 (m, 2H, Ph-H), 7.81 (m, 2H, Ph-H), 8.51 (d, 1H, J = 5.0 Hz, pyrimidinyl-H), 9.82 (brs, 1H, NH).

N-{5-[2-(4-Chloro-phenylamino)-pyrimidin-4-yl]-4-methyl-thiazol-2-yl}-3-(2-morpholin-4-yl-ethylamino)-propionamide [37]. By treatment of 3-bromo-*N*-{5-[2-(4-chloro-phenylamino)-pyrimidin-4-yl]-4-methyl-thiazol-2-yl}-propionamide with 2-morpholin-4-yl-ethylamine (or 3-amino-*N*-{5-[2-(4-chloro-phenylamino)-pyrimidin-4-yl]-4-methyl-thiazol-2-yl}-propionamide with 4-(2-chloro-ethyl)-morpholine. Anal. RP-HPLC: t_R = 11.5 min (10 – 70 % MeCN, purity > 97 %). $^1\text{H-NMR}$ (DMSO-D_6) δ : 1.52 (d, 2H, J = 7.0Hz, CH_2), 2.48 (m, 2H, CH_2), 2.52 (s, 3H, CH_3), 3.05-3.11 (m, 4H, CH_2), 3.25-3.28 (m, 6H, CH_2), 4.08 (m, 2H, CH_2), 7.13 (d, 1H, J = 5.5Hz, pyrimidinyl-H), 7.29 (m, 2H, Ph-H), 7.82 (m, 2H, Ph-H), 8.51 (d, 1H, J = 5.5 Hz, pyrimidinyl-H), 9.82 (brs, 1H, NH).

N-{5-[2-(4-Methoxy-phenylamino)-pyrimidin-4-yl]-4-methyl-thiazol-2-yl}-3-morpholin-4-yl-propionamide [38]. By treatment of 3-bromo-*N*-{5-[2-(4-methoxy-phenylamino)-pyrimidin-4-yl]-4-methyl-thiazol-2-yl}-propionamide with morpholine. Anal. RP-HPLC:

63

$t_R = 12.7$ min (0 – 60 % MeCN, purity > 90 %). $^1\text{H-NMR}$ (CDCl_3) δ : 1.20 (m, 2H, CH_2), 2.54-2.63 (m, 7H, CH_3 and CH_2), 3.34 (m, 2H, CH_2), 3.80 (m, 5H, CH_3 and CH_2), 4.03 (m, 2H, CH_2), 6.87-6.92 (m, 3H, pyrimidinyl-H and Ph-H), 7.17 (brs, 1H, NH), 7.53 (m, 2H, Ph-H), 8.35 (m, 1H, pyrimidinyl-H), 10.33 (brs, 1H, NH).

5

N-{5-[2-(3-Methoxy-phenylamino)-pyrimidin-4-yl]-4-methyl-thiazol-2-yl}-3-morpholin-4-yl-propionamide [39]. By treatment of 3-bromo-*N*-{5-[2-(3-methoxyphenylamino)-pyrimidin-4-yl]-4-methyl-thiazol-2-yl}-propionamide with morpholine. Anal. RP-HPLC: $t_R = 13.6$ min (0 – 60 % MeCN, purity > 94 %). $^1\text{H-NMR}$ (DMSO-D_6) δ : 1.20 (m, 2H, CH_2), 2.48 (m, 4H, CH_2), 2.58 (s, 3H, CH_3), 3.41-3.57 (m, 6H, CH_2), 3.79 (s, 3H, CH_3), 6.52 (d, 1H, $J = 7.0$ Hz, Ph-H), 7.07 (d, 1H, $J = 5.4$ Hz, pyrimidinyl-H), 7.15 (t, 1H, $J = 7.3$ Hz, Ph-H), 7.22 (m, 1H, Ph-H), 7.62 (brs, 1H, Ph-H), 8.46 (d, 1H, $J = 5.5$ Hz, pyrimidinyl-H), 9.58 (brs, 1H, NH). MS (ESI^+) m/z 456.01 $[\text{M}+\text{H}]^+$ ($\text{C}_{22}\text{H}_{26}\text{N}_6\text{SO}_3$ requires 454.55).

15

N-{5-[2-(4-Methoxy-phenylamino)-pyrimidin-4-yl]-4-methyl-thiazol-2-yl}-3-(4-methyl-piperazin-1-yl)-propionamide [40]. By treatment of 3-bromo-*N*-{5-[2-(4-methoxyphenylamino)-pyrimidin-4-yl]-4-methyl-thiazol-2-yl}-propionamide with 1-methyl-piperazine. Anal. RP-HPLC: $t_R = 13.0$ min (0 – 60 % MeCN, purity > 94 %). $^1\text{H-NMR}$ (DMSO-D_6) δ : 1.17 (m, 2H, CH_2), 2.11 (m, 2H, CH_2), 2.48 (m, 4H, CH_2), 2.57 (s, 3H, CH_3), 3.28 (s, 3H, CH_3), 3.30 (m, 4H, CH_2), 3.72 (s, 3H, CH_3), 6.85 (m, 2H, Ph-H), 6.97 (m, 1H, pyrimidinyl-H), 7.63 (m, 2H, Ph-H), 8.38 (d, 1H, $J = 5.1$ Hz, pyrimidinyl-H). MS (ESI^+) m/z 468.57 $[\text{M}+\text{H}]^+$ ($\text{C}_{23}\text{H}_{26}\text{N}_7\text{SO}_2$ requires 467.59).

20

N-{4-[4-(2,4-Dimethyl-thiazol-5-yl)-pyrimidin-2-ylamino]-phenyl}-acetamide [45]. By condensation between 3-dimethylamino-1-(2,4-dimethyl-thiazol-5-yl)-propanone and *N*-(4-acetamidophenyl)-guanidine nitrate. Pale solid. Mp. 219-220 °C. $^1\text{H-NMR}$ (DMSO-D_6) δ : 2.00 (s, 3H, CH_3), 2.61 (s, 3H, CH_3), 2.64 (s, 3H, CH_3), 7.04 (d, 1H, $J = 4.9$ Hz, pyrimidinyl-H), 7.48 (d, 2H, $J = 8.8$ Hz, Ph-H), 7.64 (d, 2H, $J = 8.8$ Hz, Ph-H), 8.47 (d, 1H,

25

64

$J = 5.4\text{Hz}$, pyrimidinyl-H), 9.58 (s, 1H, NH), 9.82 (s, 1H, NH). MS (ESI⁺) m/z 340.02 [M+H]⁺ (C₁₇H₁₇N₅OS requires 339.42).

N-{4-[4-(2,4-Dimethyl-thiazol-5-yl)-pyrimidin-2-ylamino]-phenyl}-*N*-methyl-acetamide

- 5 [48]. By condensation reaction of 3-dimethylamino-1-(2,4-dimethyl-thiazol-5-yl)-propanone and *N*-(4-*N*-methylacetamidophenyl)-guanidine nitrate. ¹H-NMR (DMSO-D₆) δ : 1.75 (s, 3H, CH₃), 2.62 (s, 3H, CH₃), 2.64 (s, 3H, CH₃), 3.11 (s, 3H, CH₃), 7.11 (d, 1H, $J = 4.9\text{Hz}$, pyrimidinyl-H), 7.24 (d, 2H, $J = 7.8\text{Hz}$, Ph-H), 7.83 (d, 2H, $J = 7.8\text{Hz}$, Ph-H), 8.53 (d, 1H, $J = 4.9\text{Hz}$, pyrimidinyl-H), 9.83 (brs, 1H, NH). MS (ESI⁺) m/z 354.88
- 10 [M+H]⁺ (C₁₈H₁₉N₅OS requires 353.44).

1-(4-{4-[4-(2,4-Dimethyl-thiazol-5-yl)-pyrimidin-2-ylamino]-phenyl}-piperazin-1-yl)-propan-2-ol [49].

- By treatment of [4-(2,4-dimethyl-thiazol-5-yl)-pyrimidin-2-yl]-(4-piperazin-1-yl-phenyl)-amine with 1-chloro-2-propanol. ¹H-NMR (CDCl₃) δ : 1.10 (d, 3H, $J = 6.0\text{Hz}$, CH₃), 2.26-2.33 (m, 2H, CH₂), 2.49-2.53 (m, 2H, CH₂), 2.61 (s, 3H, CH₃), 2.63 (s, 3H, CH₃), 2.76-2.80 (m, 2H, CH₂), 3.07-3.13 (m, 4H, CH₂), 3.83 (m, 1H, CH), 6.81 (d, 1H, $J = 4.4\text{Hz}$, pyrimidinyl-H), 6.88 (d, 2H, $J = 8.8\text{Hz}$, Ph-H), 7.02 (brs, 1H, OH), 7.44 (d, 2H, $J = 8.8\text{Hz}$, Ph-H), 8.30 (d, 1H, $J = 4.4\text{Hz}$, pyrimidinyl-H).
- 15

- 20 2-Chloro-*N*-{4-[4-(2,4-dimethyl-thiazol-5-yl)-pyrimidin-2-ylamino]-phenyl}-acetamide

- [50]. By treatment of [4-(2,4-dimethylthiazol-5-yl)-pyrimidin-2-yl]-(4-aminophenyl)-amine with chloroacetyl chloride. Mp. 217-219 °C. ¹H-NMR (DMSO-D₆) δ : 2.61 (s, 3H, CH₃), 2.64 (s, 3H, CH₃), 4.22 (s, 2H, CH₂), 7.05 (d, 1H, $J = 5.4\text{Hz}$, pyrimidinyl-H), 7.50 (d, 2H, $J = 8.8\text{Hz}$, Ph-H), 7.70 (d, 2H, $J = 8.8\text{Hz}$, Ph-H), 8.49 (d, 1H, $J = 4.9\text{Hz}$, pyrimidinyl-H), 9.65 (s, 1H, NH), 10.20 (s, 1H, NH). MS (ESI⁺) m/z 374.47 [M+H]⁺ (C₁₇H₁₆ClN₅OS requires 373.86).
- 25

N-{4-[4-(2,4-Dimethyl-thiazol-5-yl)-pyrimidin-2-ylamino]-phenyl}-2-morpholin-4-yl-acetamide [51].

- By treatment of 2-chloro-*N*-{4-[4-(2,4-dimethyl-thiazol-5-yl)-pyrimidin-2-ylamino]-phenyl}-acetamide with morpholine. ¹H-NMR (DMSO-D₆) δ : 2.61 (s, 3H, CH₃),
- 30

65

2.64 (s, 3H, CH₃), 3.09 (s, 2H, CH₂), 3.27-3.31 (m, 4H, CH₂), 3.62-3.64 (m, 4H, CH₂), 7.04 (d, 1H, *J* = 5.4Hz, pyrimidinyl-H), 7.54 (d, 2H, *J* = 8.8Hz, Ph-H), 7.67 (d, 2H, *J* = 8.8Hz, Ph-H), 8.48 (d, 1H, *J* = 4.9Hz, pyrimidinyl-H), 9.59 (s, 1H, NH). MS (ESI⁺) *m/z* 425.01 [M+H]⁺ (C₂₁H₂₄N₆O₂S requires 424.52).

5

N-{4-[4-(2,4-Dimethyl-thiazol-5-yl)-pyrimidin-2-ylamino]-phenyl}-2-[1,2,4]triazol-1-yl-acetamide [52]. By treatment of 2-chloro-N-{4-[4-(2,4-dimethyl-thiazol-5-yl)-pyrimidin-2-ylamino]-phenyl}-acetamide with 1*H*-[1,2,4]triazole. ¹H-NMR (DMSO-D₆) δ: 2.61 (s, 3H, CH₃), 2.63 (s, 3H, CH₃), 5.10 (s, 2H, CH₂), 7.05 (d, 1H, *J* = 5.4Hz, pyrimidinyl-H), 7.50 (d, 2H, *J* = 8.3Hz, pyrimidinyl-H), 7.69 (d, 2H, *J* = 8.9Hz, Ph-H), 7.98 (s, 1H, Aryl-H), 8.48 (d, 1H, *J* = 4.9Hz, pyrimidinyl-H), 8.53 (s, 1H, Aryl-H), 9.62 (brs, 1H, NH), 10.30 (brs, 1H, NH). MS (ESI⁺) *m/z* 406.97 (C₁₉H₁₈N₈OS requires 406.47).

N-{4-[4-(2,4-Dimethyl-thiazol-5-yl)-pyrimidin-2-ylamino]-phenyl}-2-pyrrolidin-1-yl-acetamide [53]. By treatment of 2-chloro-N-{4-[4-(2,4-dimethyl-thiazol-5-yl)-pyrimidin-2-ylamino]-phenyl}-acetamide with pyrrolidine. ¹H-NMR (DMSO-D₆) δ: 1.74 (m, 4H, CH₂), 2.58 (m, 4H, CH₂), 2.61 (s, 3H, CH₃), 2.64 (s, 3H, CH₃), 3.20 (s, 2H, CH₂), 7.04 (d, 1H, *J* = 5.4Hz, pyrimidinyl-H), 7.55 (d, 2H, *J* = 8.8Hz, Ph-H), 7.66 (d, 2H, *J* = 8.8Hz, Ph-H), 8.48 (d, 1H, *J* = 5.4Hz, pyrimidinyl-H), 9.55 (s, 1H, NH), 9.58 (s, 1H, NH). MS (ESI⁺) *m/z* 406.97 [M+H]⁺ (C₂₁H₂₄N₆OS requires 408.52).

N-{4-[4-(2,4-Dimethyl-thiazol-5-yl)-pyrimidin-2-ylamino]-phenyl}-2-imidazol-1-yl-acetamide [54]. By treatment of 2-chloro-N-{4-[4-(2,4-dimethyl-thiazol-5-yl)-pyrimidin-2-ylamino]-phenyl}-acetamide with 1*H*-imidazole. ¹H-NMR (DMSO-D₆) δ: 2.61 (s, 3H, CH₃), 2.63 (s, 3H, CH₃), 4.86 (s, 2H, CH₂), 6.88 (s, 1H, Aryl-H), 7.04 (d, 1H, *J* = 5.9Hz, pyrimidinyl-H), 7.15 (s, 1H, Aryl-H), 7.50 (d, 2H, *J* = 7.8Hz, Ph-H), 7.62 (s, 1H, Aryl-H), 7.68 (d, 2H, *J* = 7.8Hz, Ph-H), 8.48 (d, 1H, *J* = 5.4Hz, pyrimidinyl-H), 9.60 (s, 1H, NH), 10.18 (brs, 1H, NH). MS (ESI⁺) *m/z* 406.02 [M+H]⁺ (C₂₀H₁₉N₇OS requires 405.48).

3-[4-(4-Methyl-2-morpholin-4-yl-thiazol-5-yl)-pyrimidin-2-ylamino]-phenol [55]. A solution of morpholine-4-carbonitrile (10 g, 89.19 mmol) in EtOH (65 mL) was cooled on an ice bath. Anhydrous NH₃ was bubbled through the solution for 5 min, followed by hydrogen sulfide. Soon after the introduction of H₂S a white precipitate was observed. After the addition of both gases for 45 min NH₃ addition was stopped and H₂S continued for a further 15 minutes. The resulting precipitate was collected, washed with cold water, MeOH and dried under high vacuum to afford morpholine-4-carbothioic acid amide (12.83 g). Mp. 173-174 °C. ¹H-NMR (DMSO-D₆) δ: 3.54 (t, 4H, *J* = 4.9Hz, CH₂), 3.70 (m, 4H, CH₂), 7.46 (brs, 2H, NH₂). This was converted first to 1-(4-methyl-2-morpholin-4-yl-thiazol-5-yl)-ethanone with 3-bromo-pentane-2,4-dione, then with dimethoxymethyl-dimethyl-amine to 3-dimethylamino-1-(4-methyl-2-morpholin-4-yl-thiazol-5-yl)-propenone in the usual manner. The latter enaminone was condensed with *N*-(3-hydroxy-phenyl)-guanidine nitrate to afford the title compound as a pale solid. Mp. 227-229 C. ¹H-NMR (DMSO-D₆) δ: 2.49 (s, 3H, CH₃), 3.46 (t, 4H, *J* = 4.4Hz, CH₂), 3.71 (t, 4H, *J* = 4.4Hz, pyrimidinyl-H), 6.34 (d, 1H, *J* = 8.8Hz, Ph-H), 6.91 (d, 1H, *J* = 5.4Hz, pyrimidinyl-H), 7.03 (t, 1H, *J* = 7.8Hz, Ph-H), 7.20-7.22 (m, 2H, Ph-H), 8.34 (d, 1H, *J* = 5.4Hz, pyrimidinyl-H), 9.17 (s, 1H, OH/NH), 9.32 (s, 1H, NH/OH). MS (ESI⁺) *m/z* 370.10 [M+H]⁺ (C₁₈H₁₉N₅O₂S requires 369.44).

[4-(4-Methyl-2-morpholin-4-yl-thiazol-5-yl)-pyrimidin-2-yl]-(4-morpholin-4-yl-phenyl)-amine [56]. By treatment of 3-dimethylamino-1-(4-methyl-2-morpholino-thiazol-5-yl)-propenone and *N*-(4-morpholinophenyl)-guanidine nitrate. Pale solid. Mp. 229-231 °C. ¹H-NMR (DMSO-D₆) δ: 2.48 (s, 3H, CH₃), 3.02 (t, 4H, *J* = 4.0Hz, CH₂), 3.46 (t, 4H, *J* = 4.0Hz, CH₂), 3.70-3.73 (m, 8H, CH₂), 6.86 (d, 1H, *J* = 5.4Hz, pyrimidinyl-H), 6.89 (d, 2H, *J* = 9.3Hz, Ph-H), 7.60 (d, 2H, *J* = 8.8Hz, Ph-H), 8.30 (d, 1H, *J* = 5.4Hz, pyrimidinyl-H), 9.22 (s, 1H, NH). MS (ESI⁺) *m/z* 439.03 [M+H]⁺ (C₂₂H₂₆N₆O₂S requires 438.55).

N,N-Dimethyl-*N'*-[4-(4-methyl-2-morpholin-4-yl-thiazol-5-yl)-pyrimidin-2-yl]-benzene-1,4-diamine [57]. By treatment of 3-dimethylamino-1-(4-methyl-2-morpholino-thiazol-5-yl)-propenone and *N*-(4-*N,N*-dimethylaminophenyl)-guanidine nitrate. Yellow solid. ¹H-

67

NMR (DMSO- D_6) δ : 2.48 (s, 3H, CH_3), 2.82 (s, 6H, CH_3), 3.46 (t, 4H, $J = 4.9$ Hz, pyrimidinyl-H), 3.70 (t, 4H, $J = 4.9$ Hz, CH_2), 6.70 (d, 2H, $J = 8.8$ Hz, Ph-H), 6.82 (d, 1H, $J = 5.4$ Hz, pyrimidinyl-H), 7.53 (d, 2H, $J = 8.8$ Hz, Ph-H), 8.27 (d, 1H, $J = 5.4$ Hz, pyrimidinyl-H), 9.09 (s, 1H, NH). MS (ESI $^+$) m/z 397.03 $[M+H]^+$ ($C_{20}H_{24}N_6OS$ requires 396.51).

2-{4-[4-(4-Methyl-2-methylamino-thiazol-5-yl)-pyrimidin-2-ylamino]-phenyl}-ethanol [59]. By condensation between 3-dimethylamino-1-(4-methyl-2-methylamino-thiazol-5-yl)-propenone and *N*-[4-(2-hydroxy-ethyl)-phenyl]-guanidine nitrate. Pale solid. Mp 216-218 °C. Anal. RP-HPLC: $t_R = 9.1$ min (10 – 70 % MeCN, purity > 95 %). 1H -NMR (DMSO- d_6) δ : 2.46 (s, 3H, CH_3), 3.05 (s, 3H, CH_3), 3.55 (m, 2H, CH_2), 4.58 (m, 2H, CH_2), 6.85 (d, 1H, $J = 5.5$ Hz, pyrimidinyl-H), 7.08 (m, 2H, Ph-H), 7.64 (m, 2H, Ph-H), 8.01 (m, 1H, OH), 8.29 (d, 1H, $J = 5.5$ Hz, pyrimidinyl-H), 9.30 (brs, 1H, NH). MS (ESI $^+$) m/z 363.99 $[M+H]^+Na$ ($C_{17}H_{19}N_5SONa$ requires 364.43).

15

1-(4-{4-[4-(4-Methyl-2-methylamino-thiazol-5-yl)-pyrimidin-2-ylamino]-phenyl}-piperazin-1-yl)-ethanone [61]. By condensation between 3-dimethylamino-1-(4-methyl-2-methylamino-thiazol-5-yl)-propenone and *N*-[4-(4-acetyl-piperazin-1-yl)-phenyl]-guanidine nitrate. Yellow solid. Mp 213-214 °C. Anal. RP-HPLC: $t_R = 8.8$ min (10 – 70 % MeCN, purity > 95 %). 1H -NMR (DMSO- d_6) δ : 2.43 (s, 3H, CH_3), 3.02 (s, 3H, CH_3), 3.23 (s, 3H, CH_3), 2.99 (m, 2H, CH_2), 3.06 (t, 2H, CH_2), 3.57 (t, 4H, CH_2), 6.82 (d, 1H, $J = 6.0$ Hz, pyrimidinyl-H), 6.89 (d, 2H, $J = 9.0$ Hz, Ph-H), 7.62 (d, 2H, $J = 9.5$ Hz, Ph-H), 8.26 (d, 1H, $J = 5.5$ Hz, pyrimidinyl-H), 9.18 (s, 1H, NH). MS (ESI $^+$) m/z 424.07 $[M+H]^+$ ($C_{21}H_{25}N_7OS$ requires 423.54).

25

[4-(4-Methyl-2-methylamino-thiazol-5-yl)-pyrimidin-2-yl]-(4-piperazin-1-yl-phenyl)-amine [62]. By hydrolysis of 1-(4-{4-[4-(4-Methyl-2-methylamino-thiazol-5-yl)-pyrimidin-2-ylamino]-phenyl}-piperazin-1-yl)-ethanone in 2 M aq HCl. Yellow solid. Anal. RP-HPLC: $t_R = 8.8$ min (10 – 70 % MeCN, purity > 95 %). 1H -NMR (DMSO- d_6) δ : 2.45 (s, 3H, CH_3), 2.83 (t, 4H, $J = 5.9$ Hz, CH_2), 2.85 (d, 3H, $J = 4.9$ Hz, CH_2), 2.95 (t, 4H, $J = 4.9$ Hz, CH_2),

30

68

6.81 (d, 1H, $J = 5.4$ Hz, pyrimidinyl-H), 6.85 (d, 2H, $J = 9.3$ Hz, Ph-H), 7.58 (d, 2H, $J = 8.8$ Hz, Ph-H), 7.99 (m, 1H, NH), 8.26 (d, 1H, $J = 5.4$ Hz, pyrimidinyl-H), 9.14 (brs, 1H).

N-{3-[4-(4-Methyl-2-methylamino-thiazol-5-yl)-pyrimidin-2-ylamino]-benzyl}-acetamide

- 5 [63]. By condensation of 3-dimethylamino-1-(2-methylamino-4-methyl-thiazol-5-yl)-propanone and *N*-(3-guanidino-benzyl)-acetamide nitrate. Yellow solid. Mp 253-255 °C. Anal. RP-HPLC: $t_R = 11.3$ min (10 – 70 % MeCN, purity > 95 %): ^1H NMR (DMSO): δ 1.86 (s, 3H, CH₃), 2.46 (s, 3H, CH₃), 2.85 (s, 2H, CH₂), 3.09 (s, 3H, CH₃), 6.82 (d, 1H, $J = 8.0$ Hz, ph-H), 6.88 (d, 1H, $J = 5.5$ Hz, pyrimidinyl-H), 7.2 (t, 1H, $J = 8.0$ Hz, Ph-H), 7.60 (d, 1H, $J = 8.0$ Hz, Ph-H), 7.73 (s, 1H, Ph-H), and 8.32 (d, 1H, $J = 5.5$ Hz, pyrimidinyl-H). MS (ESI⁺) m/z 391.55 [M+Na] (C₁₈H₂₀N₆OSNa requires 391.46).

N-{3-[4-(2-Ethylamino-4-methyl-thiazol-5-yl)-pyrimidin-2-ylamino]-benzyl}-acetamide

- 15 [64]. By condensation of 3-dimethylamino-1-(2-ethylamino-4-methyl-thiazol-5-yl)-propanone and *N*-(3-guanidino-benzyl)-acetamide nitrate. Yellow solid. Anal. RP-HPLC: $t_R = 12.7$ min (0 – 60 % MeCN, purity > 95 %). ^1H -NMR (CD₃OD) δ : 1.17 (t, 3H, $J = 7.5$ Hz, CH₃), 1.98 (s, 3H, CH₃), 2.51 (s, 3H, CH₃), 3.36 (q, 2H, $J = 7.1$ Hz, CH₂), 4.39 (s, 2H, CH₂), 6.92 (m, 2H, pyrimidinyl-H and Ph-H), 7.25 (t, 1H, $J = 7.6$ Hz, Ph-H), 7.49 (m, 1H, Ph-H), 7.79 (sbr, 1H, Ph-H), 8.25 (d, 1H, $J = 5.5$ Hz, pyrimidinyl-H). MS (ESI⁺) m/z 383.46 [M+H]⁺ (C₁₉H₂₂N₆OS requires 382.48).

(3-Aminomethyl-phenyl)-[4-(2-ethylamino-4-methyl-thiazol-5-yl)-pyrimidin-2-yl]-amine

- 25 [65]. By hydrolysis of *N*-{3-[4-(2-ethylamino-4-methyl-thiazol-5-yl)-pyrimidin-2-ylamino]-benzyl}-acetamide. Yellow solid. Mp 290-292 °C. Anal. RP-HPLC: $t_R = 10.6$ min (10 – 70 % MeCN, purity > 95 %). ^1H -NMR (DMSO-D₆) δ : 1.31 (t, 3H, $J = 7.0$ Hz, CH₃), 2.64 (s, 3H, CH₃), 3.54 (m, 2H, CH₂), 4.11 (m, 2H, CH₂), 7.14 (d, 1H, $J = 6.0$ Hz, pyrimidinyl-H), 7.22 (d, 1H, $J = 8.0$ Hz, Ph-H), 7.45 (t, 1H, $J = 8.0$ Hz, Ph-H), 7.74 (d, 1H, $J = 8.0$ Hz, Ph-H), 7.95 (s, 1H, Ph-H), 8.53 (d, 1H, $J = 6.0$ Hz, pyrimidinyl-H). MS (ESI⁺) m/z 341.20 [M+H]⁺ (C₁₇H₂₀N₆S requires 340.45).

N-{3-[4-(2,4-Dimethyl-thiazol-5-yl)-pyrimidin-2-ylamino]-benzyl}-acetamide [66]. By treatment of 3-dimethylamino-1-(2,4-dimethyl-thiazol-5-yl)-propenone and *N*-(3-guanidino-benzyl)-acetamide nitrate. Yellow solid. Mp 206-207 °C. Anal. RP-HPLC: t_R = 13.6 min (0 – 60 % MeCN, purity > 95 %). ¹H-NMR (DMSO-D₆) δ : 1.99 (s, 3H, CH₃),
 5 2.65 (s, 3H, CH₃), 2.68 (s, 3H, CH₃), 4.38 (s, 2H, CH₂), 6.94 (d, 1H, J = 7.5 Hz, Ph-H), 7.01 (d, 1H, J = 5.5 Hz, pyrimidinyl-H), 7.26 (d, 1H, J = 8.0 Hz, Ph-H), 7.56 (d, 1H, J = 7.5 Hz, Ph-H), 7.70 (s, 1H, Ph-H), 8.40 (d, 1H, J = 5.5 Hz, pyrimidinyl-H). MS (ESI⁺) m/z 345.42 [M+H]⁺ (C₁₈H₁₉N₅OS requires 353.44).

10 (3-Aminomethyl-phenyl)-[4-(4-methyl-2-methylamino-thiazol-5-yl)-pyrimidin-2-yl]-amine [67]. By treatment of *N*-{3-[4-(4-methyl-2-methylamino-thiazol-5-yl)-pyrimidin-2-ylamino]-benzyl}-acetamide with HCl/MeOH. Yellow solid. Mp 287-289 °C. Anal. RP-HPLC: t_R = 10.1 min (0 – 60 % MeCN, purity 92 %). ¹H-NMR (DMSO-D₆) δ : 2.68 (s, 3H, CH₃), 3.17 (s, 3H, CH₃), 4.17 (m, 2H, CH₂), 7.23 (d, 1H, J = 6.0 Hz, pyrimidinyl-H), 7.27
 15 (d, 1H, J = 8.0 Hz, Ph-H), 7.50 (t, 1H, J = 8.0 Hz, Ph-H), 7.70 (d, 1H, J = 7.5 Hz, Ph-H), 7.75 (s, 1H, Ph-H), 8.44 (d, 1H, J = 6.0 Hz, pyrimidinyl-H). MS (ESI⁺) m/z 327.37 [M+H]⁺ (C₁₆H₁₈N₆S requires 326.42).

{3-[4-(2-Ethylamino-4-methyl-thiazol-5-yl)-pyrimidin-2-ylamino]-phenyl}-methanol [68].
 20 By condensation of 3-dimethylamino-1-(2-ethylamino-4-methyl-thiazol-5-yl)-propenone and *N*-(3-hydroxymethyl-phenyl)-guanidine nitrate. Yellow solid. Anal. RP-HPLC: t_R = 12.6 min (0 – 60 % MeCN, purity > 95 %). ¹H-NMR (CD₃OD) δ : 1.28 (t, 3H, J = 7.3 Hz, CH₃), 2.52 (s, 3H, CH₃), 3.35 (q, 2H, J = 7.1 Hz, CH₂), 4.63 (s, 2H, CH₂), 5.49 (d, 1H, J = 5.5 Hz, pyrimidinyl-H), 7.63 (d, 1H, J = 7.6 Hz, Ph-H), 7.27 (t, 1H, J = 7.9 Hz, Ph-H),
 25 7.52 (m, 1H, Ph-H), 7.79 (sbr, 1H, Ph-H), 8.26 (d, 1H, J = 5.5 Hz, pyrimidinyl-H). MS (ESI⁺) m/z 346.44 [M+H]⁺ (C₁₇H₁₉N₅OS requires 341.43).

[4-(4-Methyl-2-morpholin-4-yl-thiazol-5-yl)-pyrimidin-2-yl]-(3-nitro-phenyl)-amine [71].
 By condensation of 3-dimethylamino-1-(4-methyl 2-morpholin-4-yl-thiazol-5-yl)-
 30 propenone and *N*-(3-nitro-phenyl)-guanidine nitrate. Yellow solid. Anal. RP-HPLC: t_R =

70

16.7 min (10 – 70 % MeCN, purity > 95 %). ¹H-NMR (DMSO-D₆) δ: 2.51 (s, 3H, CH₃), 3.50 (t, 4H, *J* = 4.5 Hz, CH₂), 3.72 (t, 4H, *J* = 4.5 Hz, CH₂), 7.06 (d, 1H, *J* = 5.5 Hz, pyrimidinyl-H), 7.55 (t, 1H, *J* = 8.5 Hz, Ph-H), 7.77 (d, 1H, *J* = 8.5 Hz, Ph-H), 7.97 (d, 1H, *J* = 8.5 Hz, Ph-H), 8.44 (d, 1H, *J* = 5.5 Hz, pyrimidinyl-H), 9.03 (s, 1H, Ph-H), 10.06 (sbr, 1H, NH). MS (ESI⁺) *m/z* 399.20 [M+H]⁺ (C₁₈H₁₈N₆O₃S requires 398.44).

{2-Chloro-5-[4-(2,4-dimethyl-thiazol-5-yl)-pyrimidin-2-ylamino]-phenyl}-methanol [72].

By condensation of 3-dimethylamino-1-(2,4-dimethyl-thiazol-5-yl)-propenone and *N*-(4-chloro-3-hydroxymethyl-phenyl)-guanidine nitrate. Yellow solid. Mp 245-246 °C. Anal.

10 RP-HPLC: *t_R* = 14.6 min (10 – 70 % MeCN, purity > 95 %). ¹H-NMR (DMSO-D₆) δ: 2.62 (s, 3H, CH₃), 2.63 (s, 3H, CH₃), 4.53 (d, 2H, *J* = 5.5 Hz, CH₂), 5.34 (m, 1H, OH), 7.08 (d, 1H, *J* = 5.5 Hz, pyrimidinyl-H), 7.29 (m, 1H, Ph-H), 7.73 (m, 1H, Ph-H), 7.94 (s, 1H, Ph-H), 8.51 (d, 1H, *J* = 5.5 Hz, pyrimidinyl-H), 9.78 (s, 1H, NH). MS (ESI⁺) *m/z* 347.11 [M+H]⁺ (C₁₆H₁₅ClN₄OS requires 346.84).

15

{2-Chloro-5-[4-(4-methyl-2-methylamino-thiazol-5-yl)-pyrimidin-2-ylamino]-phenyl}-methanol [73].

By condensation of 3-dimethylamino-1-(2-methylamino-4-methyl-thiazol-5-yl)-propenone and *N*-(4-chloro-3-hydroxymethyl-phenyl)-guanidine nitrate. Yellow solid. Mp 191-193 °C. Anal. RP-HPLC: *t_R* = 11.5 min (10 – 70 % MeCN, purity > 90 %). ¹H-

20 NMR (DMSO-D₆) δ: 2.46 (s, 3H, CH₃), 3.08 (s, 3H, CH₃), 4.52 (d, 2H, *J* = 6.0 Hz, CH₂), 5.29 (m, 1H, OH), 6.89 (d, 1H, *J* = 5.5 Hz, pyrimidinyl-H), 7.25 (d, 1H, *J* = 8.5 Hz, Ph-H), 7.73 (m, 1H, Ph-H), 7.94 (s, 1H, Ph-H), 8.03 (sbr, 1H, NH), 8.32 (d, 1H, *J* = 5.5 Hz, pyrimidinyl-H), 9.55 (s, 1H, NH). MS (ESI⁺) *m/z* 361.89 [M] (C₁₆H₁₆ClN₅OS requires 361.85).

25

[4-(2-Amino-4-methyl-thiazol-5-yl)-pyrimidin-2-yl]-(4-methanesulfonyl-phenyl)-amine

[76]. By condensation of *N*-[5-(3-dimethylamino-acryloyl)-4-methyl-thiazol-2-yl]-*N,N*-dimethyl-formamide and *N*-(4-methanesulfonyl-phenyl)-guanidine nitrate. Yellow solid.

Anal. RP-HPLC: *t_R* = 13.2 min (0 – 60 % MeCN, purity > 97 %). ¹H-NMR (DMSO-d₆) δ:

30 1.98 (s, 3H, CH₃), 2.54 (s, 3H, CH₃), 7.06 (d, 1H, *J* = 5.5 Hz, pyrimidinyl-H), 7.80 (m, 2H,

71

Ph-H), 8.00 (m, 2H, Ph-H), 8.45 (d, 1H, $J = 5.5$ Hz, pyrimidinyl-H), 10.05 (sbr, 2H, NH_2). MS (ESI^+) m/z 362.38 $[\text{M}+\text{H}]^+$ ($\text{C}_{15}\text{H}_{15}\text{N}_5\text{O}_2\text{S}_2$ requires 361.44).

[4-(2-Methoxy-ethoxy)-3-nitro-phenyl]-[4-(4-methyl-2-methylamino-thiazol-5-yl)-

- 5 *pyrimidin-2-yl]-amine [77].* By alkylation of [4-(2-aminomethyl-4-methyl-thiazol-5-yl)-pyrimidin-2-yl]-(4-fluoro-3-nitrophenyl)-amine with 2-methoxy-ethanol. Yellow solid. Anal. RP-HPLC: $t_R = 12.8$ min (10 – 70 % MeCN, purity > 90 %). $^1\text{H-NMR}$ (DMSO-D_6) δ : 2.41 (s, 3H, CH_3), 2.86 (d, 3H, $J = 4.5$ Hz, CH_3), 3.24 (s, 3H, CH_3), 3.66 (m, 2H, CH_2), 4.23 (m, 2H, CH_2), 6.93 (d, 1H, $J = 6.0$ Hz, pyrimidinyl-H), 7.32 (d, 1H, $J = 5.0$ Hz, Ph-H), 7.79 (m, 1H, Ph-H), 8.08 (d, 1H, $J = 5.0$ Hz, Ph-H), 8.35 (d, 1H, $J = 5.0$ Hz, pyrimidinyl-H), 8.54 (m, 1H, NH), 9.68 (s, 1H, NH). MS (ESI^+) m/z 417.08 $[\text{M}+\text{H}]^+$ ($\text{C}_{18}\text{H}_{20}\text{N}_6\text{O}_4\text{S}$ requires 416.46).

[4-(2,4-Dimethyl-thiazol-5-yl)-pyrimidin-2-yl]-[3-(2-morpholin-4-yl-ethoxymethyl)-

- 15 *phenyl]-amine [81].* By alkylation of {3-[4-(2,4-dimethyl-thiazol-5-yl)-pyrimidin-2-ylamino]-phenyl}-methanol with 4-(2-chloro-ethyl)-morpholine. Yellow solid. Anal. RP-HPLC: $t_R = 8.5$ min (10 – 70 % MeCN, purity > 95 %). $^1\text{H-NMR}$ (DMSO-D_6) δ : 2.56 (s, 3H, CH_3), 2.57 (s, 3H, CH_3), 2.63 (m, 2H, CH_2), 3.58 (m, 4H, $J = 4.5$ Hz, $\text{CH}_2 \times 2$), 4.14 (t, 2H, $J = 7.5$ Hz, CH_2), 4.58 (d, 2H, $J = 5.0$ Hz, CH_2), 5.28 (t, 1H, $J = 5.5$ Hz, NH), 7.06 (d, 20 1H, $J = 5.5$ Hz, pyrimidinyl-H), 7.26 (m, 2H, Ph-H), 7.36 (s, 1H, Ph-H), 7.42 (m, 1H, Ph-H), 8.43 (d, 1H, $J = 5.5$ Hz, pyrimidinyl-H), 9.12 (sbr, 1H, Ph-H). MS (ESI^+) m/z 426.46 $[\text{M}+\text{H}]^+$ ($\text{C}_{22}\text{H}_{27}\text{N}_5\text{O}_2\text{S}$ requires 425.55).

C,C,C-Trifluoro-N-{3-[4-(4-methyl-2-methylamino-thiazol-5-yl)-pyrimidin-2-ylamino]-

- 25 *benzyl}-methanesulfonamide [82].* By treatment of (3-aminomethyl-phenyl)-[4-(4-methyl-2-methylamino-thiazol-5-yl)-pyrimidin-2-yl]-amine with trifluoro-methanesulfonyl chloride. Yellow solid. Anal. RP-HPLC: $t_R = 11.6$ min (0 – 60 % MeCN, purity ~ 90 %). $^1\text{H-NMR}$ (CD_3OD) δ : 2.53 (s, 3H, CH_3), 2.97 (s, 3H, CH_3), 4.39 (s, 2H, CH_2), 6.94 (d, 1H, $J = 5.5$ Hz, pyrimidinyl-H), 7.01 (d, 1H, $J = 7.5$ Hz, Ph-H), 7.31 (d, 1H, $J = 8.0$ Hz, Ph-H),

72

7.60 (d, 1H, $J = 7.5\text{Hz}$, Ph-H), 7.67 (s, 1H, Ph-H), 8.28 (d, 1H, $J = 5.5\text{ Hz}$, pyrimidinyl-H). MS (ESI⁺) m/z 459.33 [M+H]⁺ (C₁₇H₁₇F₃N₆O₂S requires 458.48).

N-{3-[4-(4-Methyl-2-methylamino-thiazol-5-yl)-pyrimidin-2-ylamino]-benzyl}-

5 *methanesulfonamide* [83]. By treatment of (3-aminomethyl-phenyl)-[4-(4-methyl-2-methylamino-thiazol-5-yl)-pyrimidin-2-yl]-amine with methane-sulfonyl chloride. Yellow solid. Anal. RP-HPLC: $t_R = 12.8\text{ min}$ (0 – 60 % MeCN, purity >95 %). ¹H-NMR (CD₃OD) δ : 2.53 (s, 3H, CH₃), 2.88 (s, 3H, CH₃), 2.98 (s, 3H, CH₃), 4.28 (s, 2H, CH₂), 6.94 (d, 1H, $J = 5.5\text{Hz}$, pyrimidinyl-H), 7.02 (d, 1H, $J = 7.5\text{Hz}$, Ph-H), 7.29 (t, 1H, $J = 8.0\text{Hz}$, Ph-H),
10 7.52 (d, 1H, $J = 8.0\text{Hz}$, Ph-H), 7.90 (s, 1H, Ph-H), 7.98 (s, 1H, NH), 8.28 (d, 1H, $J = 5.5\text{ Hz}$, pyrimidinyl-H).

[4-(2-Amino-4-methyl-thiazol-5-yl)-pyrimidin-2-yl]-(3-methanesulfonyl-phenyl)-amine

[84]. By condensation of *N*'-[5-(3-dimethylamino-acryloyl)-4-methyl-thiazol-2-yl]-*N,N*-
15 dimethyl-formamidine and *N*-(3-methanesulfonyl-phenyl)-guanidine nitrate. Yellow solid. Anal. RP-HPLC: $t_R = 13.1\text{ min}$ (0 – 60 % MeCN, purity > 97 %). ¹H-NMR (DMSO-*d*₆) δ : 2.55 (s, 3H, CH₃), 3.19 (s, 3H, CH₃), 6.97 (d, 1H, $J = 5.5\text{ Hz}$, pyrimidinyl-H), 7.47 (m, 1H, Ph-H), 7.54 (t, 1H, $J = 7.5\text{Hz}$, Ph-H), 8.08 (m, 1H, Ph-H), 8.34 (brs, 1H, Ph-H), 8.40 (d, 1H, $J = 5.5\text{ Hz}$, pyrimidinyl-H), 9.86 (sbr, 2H, NH₂). MS (ESI⁺) m/z 362.38 [M+H]⁺
20 (C₁₅H₁₅N₅O₂S₂ requires 361.44).

[4-(2,4-Dimethyl-thiazol-5-yl)-pyrimidin-2-yl]-(4-methanesulfonyl-phenyl)-amine [85]. By condensation of 3-dimethylamino-1-(2,4-dimethyl-thiazol-5-yl)-propanone and *N*-(4-methanesulfonyl-phenyl)-guanidine nitrate. Yellow solid. Anal. RP-HPLC: $t_R = 16.6\text{ min}$
25 (0 – 60 % MeCN, purity > 97 %). ¹H-NMR (DMSO-*d*₆) δ : 2.65 (s, 3H, CH₃), 2.66 (s, 3H, CH₃), 3.15 (s, 3H, CH₃), 7.22 (d, 1H, $J = 5.5\text{ Hz}$, pyrimidinyl-H), 7.84 (d, 2H, $J = 9.0\text{Hz}$, Ph-H), 8.03 (d, 2H, $J = 9.0\text{Hz}$, Ph-H), 8.61 (d, 1H, $J = 5.5\text{ Hz}$, pyrimidinyl-H), 10.23 (sbr, 1H, NH). MS (ESI⁺) m/z 361.17 [M+H]⁺ (C₁₆H₁₆N₄O₂S₂ requires 361.46).

- 3,4-Dimethyl-5-[2-(4-morpholin-4-yl-phenylamino)-pyrimidin-4-yl]-3H-thiazol-2-one [92]. Methylammonium *N*-methylthio-carbamate (13.1 g, 0.105 mol; prepared from methylamine and carbonyl sulfide as described, Y. Gelernt *et al.* 1974, *J. Chem. Soc. Perkin Trans. I*, 2610) was partially dissolved in MeOH (150 mL). 3-Chloro-pentane-2,4-dione (14.9 mL, 0.125 mol) was added drop-wise at room temperature, producing a gradual exotherm to 40 °C. After stirring at room temperature for 1 h, the solvent was removed *in vacuo*. The residue was treated with H₂O (50 mL) and was extracted with CH₂Cl₂ (3 × 50 mL). The combined organic fractions were washed (brine), dried (Na₂SO₄), filtered, and evaporated *in vacuo* to an amber-coloured oil. This was purified by chromatography (300 g SiO₂, eluting with 1:1 heptane/Et₂O to obtain non-cyclized adduct, then Et₂O to obtain 5-acetyl-3,4-dimethyl-3H-thiazol-2-one, which was recrystallized from EtOH as colourless needles (14.2 g). ¹H-NMR (CDCl₃): δ 2.34 (s, 3H), 2.59 (s, 3H), 3.33 (s, 3H). IR (ATR): 1655 and 1621 cm⁻¹ (CO str).
- 5-Acetyl-3,4-dimethyl-3H-thiazol-2-one (4.64 g, 27.10 mmol) and dimethylformamide dimethyl acetal (8.4 mL, 59.62 mmol) were mixed in a dry, argon-flushed flask, and heated at 100 °C for 3 h. The mixture was cooled, producing some precipitation, which was enhanced by the addition of an equal volume of Et₂O. The resulting orange solid was filtered and washed with Et₂O to give 2.73g of 5-(3-dimethylamino-acryloyl)-3,4-dimethyl-3H-thiazol-2-one. ¹H-NMR (*d*₆-DMSO): δ 2.52 (s, 3H), 2.82 (bs, 3H), 3.11 (bs, 3H), 3.22 (s, 3H), 5.10 (d, 1H, J=12.2Hz), 7.61 (d, 1H, J=11.7Hz). IR (ATR): 1669 and 1630 cm⁻¹ (CO str).
- Condensation between 5-(3-dimethylamino-acryloyl)-3,4-dimethyl-3H-thiazol-2-one and *N*-(4-morpholin-4-yl-phenyl)-guanidine nitrate afforded the title compound. Anal. RP-HPLC: *t*_R = 17.5 min (0 – 60 % MeCN in 0.1 % aq CF₃COOH over 20 min, 1 mL/min, purity > 95 %). ¹H-NMR (DMSO-*d*₆) δ: 2.48 (3H, s, CH₃), 3.03 (4H, m, 2 x morph-NCH₂), 3.08 (3H, s, CH₃), 3.72 (4H, m, 2 x morph-OCH₂), 6.85 (1H, d, J=5.2, pyrim-H), 6.89 (2H, d, J=9.2, 2 x ArH), 7.57 (2H, d, J=9.2, 2 x ArH), 8.36 (1H, d, J=5.2, pyrim-H) and 9.35 (1H, s, NH). MS (ESI⁺) *m/z* 384[M+H]⁺ (C₁₉H₂₁N₅O₂S requires 383.5).

3,4-Dimethyl-5-{2-[4-(4-methyl-piperazin-1-yl)-phenylamino]-pyrimidin-4-yl}-3H-thiazol-2-one [93]. This compound was prepared by condensation between 5-(3-dimethylamino-acryloyl)-3,4-dimethyl-3H-thiazol-2-one and *N*-[4-(4-methyl-piperazin-1-yl)-phenyl]-guanidine nitrate. Pale solid. Anal. RP-HPLC: t_R = 10.9 min (0 – 60 % MeCN in 0.1 % aq CF₃COOH over 20 min, 1 mL/min, purity > 97 %). ¹H-NMR (CDCl₃) δ : 2.42 (m, 4H, CH₂), 2.53 (s, 3H, CH₃), 3.04 (m, 4H, CH₂), 3.28 (s, 3H, CH₃), 6.84 (d, 1H, J = 5.5 Hz, pyrimidinyl-H), 6.86 (d, 2H, J = 9.0 Hz, Ph-H), 7.54 (d, 2H, J = 9.0 Hz, Ph-H), 8.35 (d, 1H, J = 5.5 Hz, pyrimidinyl-H), and 9.33 (s, 1H, NH).

Example 3

Kinase assays. The compounds from Example 2 above were investigated for their ability to inhibit the enzymatic activity of various protein kinases. This was achieved by measurement of incorporation of radioactive phosphate from ATP into appropriate polypeptide substrates. Recombinant protein kinases and kinase complexes were produced or obtained commercially. Assays were performed using 96-well plates and appropriate assay buffers (typically 25 mM β -glycerophosphate, 20 mM MOPS, 5 mM EGTA, 1 mM DTT, 1 mM Na₃VO₃, pH 7.4), into which were added 2 - 4 μ g of active enzyme with appropriate substrates. The reactions were initiated by addition of Mg/ATP mix (15 mM MgCl₂ + 100 μ M ATP with 30-50 kBq per well of [γ -³²P]-ATP) and mixtures incubated as required at 30 °C. Reactions were stopped on ice, followed by filtration through p81 filterplates or GF/C filterplates (Whatman Polyfiltronics, Kent, UK). After washing 3 times with 75 mM aq orthophosphoric acid, plates were dried, scintillant added and incorporated radioactivity measured in a scintillation counter (TopCount, Packard Instruments, Pangbourne, Berks, UK). Compounds for kinase assay were made up as 10 mM stocks in DMSO and diluted into 10 % DMSO in assay buffer. Data was analysed using curve-fitting software (GraphPad Prism version 3.00 for Windows, GraphPad Software, San Diego California USA) to determine IC₅₀ values (concentration of test compound which inhibits kinase activity by 50 %).

- CDK 7 and 9 assays.* CTD peptide substrate (biotinyl-Ahx-(Tyr-Ser-Pro-Thr-Ser-Pro-Ser)₄-NH₂; 1 – 2 mg/mL) and recombinant human CDK7/cyclin H, CDK9/cyclin T1, or CDK9/cyclin K (0.5 – 2 µg) were incubated for 45 min at 30 °C in the presence of varying
5 amounts of test compound in 20 mM MOPS pH 7.2, 25mM β-glycerophosphate, 5 mM EGTA, 1 mM DTT, 1mM sodium vanadate, 15 mM MgCl₂, and 100 µM ATP (containing a trace amount of ³²PγATP) in a total volume of 25 µL in a 96-well microtiter plate. The reaction was stopped by placing the plate on ice for 2 min. Avidin (50 µg) was added to each well, and the plate was incubated at room temp for 30 min. The samples were
10 transferred to a 96-well P81 filter plate, and washed (4 x 200 µL per well) with 75 mM phosphoric acid. Microscint 40 scintillation liquid (50 µL) was added to each well, and the amount of ³²P incorporation for each sample was measured using a Packard Topcount microplate scintillation counter.
- 15 *Aurora-A (human) kinase assay.* This was achieved by measurement of incorporation of radioactive phosphate from ATP into Kemptide substrate (LRRASLG), upon phosphorylation by commercially obtained aurora-A kinase. Assays were performed using 96-well plates and appropriate assay buffers (8 mM MOPS, 0.2 mM EDTA, pH 7.0), into which were added 5 – 10 ng of active enzyme with 200 µM substrate (Kemptide). The
20 reactions were initiated by addition of Mg/ATP mix (10 mM MgAcetate + 15 µM ATP with 30-50 kBq per well of [γ-³³P]-ATP) and mixtures incubated for 40 min at room temperature. Reactions were stopped by addition of 3% phosphoric acid, followed by filtration through p81 filterplates (Whatman Polyfiltronics, Kent, UK). After washing 5 times with 75 mM aq orthophosphoric acid and once in methanol, plates were dried,
25 scintillant added and incorporated radioactivity measured in a scintillation counter (TopCount, Packard Instruments, Pangbourne, Berks, UK). Compounds for kinase assay were made up as 10 mM stocks in DMSO and diluted into 10 % DMSO in assay buffer. Data was analysed using curve-fitting software (XLfit version 2.0.9, IDBS, Guildford,

Surrey, UK) to determine IC₅₀ values (concentration of test compound which inhibits kinase activity by 50 %).

Results are summarized in **Table 2** and a more extensive kinase selectivity panel for
5 selected compounds is shown in **Table 3**.

Example 4

MTT cytotoxicity assay. The compounds from **Example 2** were subjected to a standard cellular proliferation assay using human tumour cell lines obtained from the ATCC
10 (American Type Culture Collection, 10801 University Boulevard, Manassas, VA 20110-2209, USA). Standard 72-h MTT (thiazolyl blue; 3-[4,5-dimethylthiazol-2-yl]-2,5-diphenyltetrazolium bromide) assays were performed (Haselsberger, K.; Peterson, D. C.; Thomas, D. G.; Darling, J. L. *Anti Cancer Drugs* 1996, 7, 331-8; Loveland, B. E.; Johns, T. G.; Mackay, I. R.; Vaillant, F.; Wang, Z. X.; Hertzog, P. J. *Biochemistry International*
15 1992, 27, 501-10). In short: cells were seeded into 96-well plates according to doubling time and incubated overnight at 37 °C. Test compounds were made up in DMSO and a 1/3 dilution series prepared in 100 µL cell media, added to cells (in triplicates) and incubated for 72 ho at 37 °C. MTT was made up as a stock of 5 mg/mL in cell media and filter-sterilised. Media was removed from cells followed by a wash with 200 µL PBS. MTT
20 solution was then added at 20 µL per well and incubated in the dark at 37 °C for 4 h. MTT solution was removed and cells again washed with 200 µL PBS. MTT dye was solubilised with 200 µL per well of DMSO with agitation. Absorbance was read at 540 nm and data analysed using curve-fitting software (GraphPad Prism version 3.00 for Windows, GraphPad Software, San Diego California USA) to determine IC₅₀ values (concentration of
25 test compound which inhibits cell growth by 50 %). Results are summarized in **Table 4** and more extensive data for selected compounds is presented in **Table 5**.

Example 5*Anti-HIV efficacy evaluation in fresh human PBMCs*

Representative compounds of the present invention were tested for antiviral activity against HIV-1 in human peripheral blood mononuclear cells (PBMCs) using the clinical
5 paediatric HIV strain RoJo or WeJo. PBMCs were cultured under conditions which promote cell survival and HIV replication. Antiviral activity was tested for from 6 – 9 log₁₀ serial dilutions of a 100 µM compound stock solution in DMSO. The following parameters were derived: IC₅₀ and IC₉₀ (concentrations inhibiting virus replication by 50 and 90 %, respectively, TC₅₀ (concentration decreasing cell viability by 50 %), and TI (therapeutic
10 index: TC₅₀ / IC₅₀).

Fresh PBMCs, seronegative for HIV and HBV, were isolated from screened donors (Interstate Blood Bank, Inc. Memphis, TN). Cells were pelleted / washed 2-3 times by low speed centrifugation and re-suspension in PBS to remove contaminating platelets. The
15 Leukophoresed blood was then diluted with Dulbecco's Phosphate Buffered Saline (DPBS) and layered over Lymphocyte Separation Medium (LSM; Cellgro® by Mediatech, Inc.; density 1.078 ± 0.002 g/mL; Cat.# 85-072-CL) in a 50 mL centrifuge tube and then centrifuged. Banded PBMCs were gently aspirated from the resulting interface and subsequently washed with PBS by low speed centrifugation. After the final wash, cells
20 were enumerated by trypan blue exclusion and re-suspended in RPMI 1640 supplemented with fetal bovine serum (FBS), and L-glutamine, Phytohemagglutinin (PHA-P, Sigma). The cells were allowed to incubate at 37 °C. After incubation, PBMCs were centrifuged and resuspended in RPMI 1640 with FBS, L-glutamine, penicillin, streptomycin, gentamycin, and recombinant human IL-2 (R&D Systems, Inc). IL-2 is included in the
25 culture medium to maintain the cell division initiated by the PHA mitogenic stimulation. PBMCs were maintained in this with bi-weekly medium changes until used in the assay protocol. Cells were kept in culture for a maximum of two weeks before being deemed too old for use in assays and discarded. Monocytes were depleted from the culture as the result of adherence to the tissue culture flask.






For the standard PBMC assay, PHA-P stimulated cells from at least two normal donors were pooled, diluted and plated in the interior wells of a 96-well round bottom microplate. Pooling of mononuclear cells from more than one donor was used to minimise the variability observed between individual donors, which results from quantitative and qualitative differences in HIV infection and overall response to the PHA and IL-2 of primary lymphocyte populations. Each plate contained virus/cell control wells (cells plus virus), experimental wells (drug plus cells plus virus) and compound control wells (drug plus media without cells, necessary for MTS monitoring of cytotoxicity). Since HIV-1 is not cytopathic to PBMCs, this allows the use of the same assay plate for both antiviral activity and cytotoxicity measurements. Test drug dilutions were prepared in microtiter tubes and each concentration was placed in appropriate wells using the standard format. A predetermined dilution of virus stock was placed in each test well (final MOI \cong 0.1). The PBMC cultures were maintained for seven days following infection at 37 °C, 5 % CO₂. After this period, cell-free supernatant samples were collected for analysis of reverse transcriptase activity and/or HIV p24 content. Following removal of supernatant samples, compound cytotoxicity was measured by addition of MTS to the plates for determination of cell viability. Wells were also examined microscopically and any abnormalities were noted.







Reverse transcriptase activity assay: A microtiter plate-based reverse transcriptase (RT) reaction was utilised (Buckheit et al., AIDS Research and Human Retroviruses 7:295-302, 1991). Tritiated thymidine triphosphate (³H-TTP, 80 Ci/mmol, NEN) was received in 1:1 dH₂O:Ethanol at 1 mCi/mL. Poly rA:oligo dT template:primer (Pharmacia) was prepared as a stock solution, followed by aliquoting and storage at -20 °C. The RT reaction buffer was prepared fresh on a daily basis. The final reaction mixture was prepared by combining ³H-TTP, dH₂O, poly rA:oligo dT stock and reaction buffer. This reaction mixture was placed in a round bottom microtiter plate and supernatant containing virus was added and mixed. The plate was incubated at 37 °C for 60 minutes. Following incubation, the reaction volume was spotted onto DE81 filter-mats (Wallac), in a sodium phosphate buffer or 2X SSC (Life Technologies). Next they were washed in distilled water, in 70 % ethanol, and






then dried. Incorporated radioactivity (counts per minute, CPM) was quantified using standard liquid scintillation techniques. The results for selected compounds of the invention are shown below in **Table 6**.





- 5 Various modifications and variations of the described aspects of the invention will be apparent to those skilled in the art without departing from the scope and spirit of the invention. Although the invention has been described in connection with specific preferred embodiments, it should be understood that the invention as claimed should not be unduly limited to such specific embodiments. Indeed, various modifications of the described modes
- 10 of carrying out the invention which are obvious to those skilled in the relevant fields are intended to be within the scope of the following claims.





Table 1. Example compounds





No.	Structure	Name
1		[4-(2-Amino-4-methyl-thiazol-5-yl)-pyrimidin-2-yl]-(4-morpholin-4-yl-phenyl)-amine
2		[4-(2,4-Dimethyl-thiazol-5-yl)-pyrimidin-2-yl]-(4-morpholin-4-yl-phenyl)-amine
3		[4-(2-N-Methylamino-4-methyl-thiazol-5-yl)-pyrimidin-2-yl]-(4-morpholinophenyl)-amine
4		[4-(2-Ethylamino-4-methyl-thiazol-5-yl)-pyrimidin-2-yl]-(4-morpholin-4-yl-phenyl)-amine
5		1-(4-{4-[4-(2,4-Dimethyl-thiazol-5-yl)-pyrimidin-2-ylamino]-phenyl}-piperazin-1-yl)-ethanone

6		[4-(2,4-Dimethyl-thiazol-5-yl)-pyrimidin-2-yl]-(4-piperazin-1-yl)-amine
7		[4-(2,4-Dimethyl-thiazol-5-yl)-pyrimidin-2-yl]-[4-(4'-2''-ethoxyethanolpiperazino)-phenyl]-amine
8		3-(4-{4-[4-(2,4-Dimethyl-thiazol-5-yl)-pyrimidin-2-ylamino]-phenyl}-piperazin-1-yl)-propan-1-ol
9		2-(4-{4-[4-(2,4-Dimethyl-thiazol-5-yl)-pyrimidin-2-ylamino]-phenyl}-piperazin-1-yl)-ethanol
10		[4-(2,4-Dimethyl-thiazol-5-yl)-pyrimidin-2-yl]-[4-(4-methanesulfonyl-piperazin-1-yl)-phenyl]-amine
11		[4-(4-Benzyl-piperazin-1-yl)-phenyl]-[4-(4-methyl-2-methylamino-thiazol-5-yl)-pyrimidin-2-yl]-amine





12		[4-(2,4-Dimethyl-thiazol-5-yl)-pyrimidin-2-yl]-[4-(4-methyl-piperazin-1-yl)-phenyl]-amine
13		3-Amino-N-{4-methyl-5-[2-(3-nitro-phenylamino)-pyrimidin-4-yl]-thiazol-2-yl}-propionamide
14		(2S)-2-Amino-3-hydroxy-N-{4-methyl-5-[2-(3-nitro-phenylamino)-pyrimidin-4-yl]-thiazol-2-yl}-propionamide
15		(2R,3R)-2-Amino-3-hydroxy-N-{4-methyl-5-[2-(3-nitro-phenylamino)-pyrimidin-4-yl]-thiazol-2-yl}-butyramide
16		(2R)-2-Amino-N-{4-methyl-5-[2-(3-nitro-phenylamino)-pyrimidin-4-yl]-thiazol-2-yl}-butyramide





17		(2S,3S)-2-Amino-3-hydroxy-N-{4-methyl-5-[2-(3-nitro-phenylamino)-pyrimidin-4-yl]-thiazol-2-yl}-butyramide
18		4-Amino-N-{4-methyl-5-[2-(3-nitro-phenylamino)-pyrimidin-4-yl]-thiazol-2-yl}-butyramide
19		3-Amino-N-{4-methyl-5-[2-(4-morpholin-4-yl-phenylamino)-pyrimidin-4-yl]-thiazol-2-yl}-propionamide
20		3-Bromo-N-{5-[2-(4-fluoro-phenylamino)-pyrimidin-4-yl]-4-methyl-thiazol-2-yl}-propionamide

21		N-{5-[2-(4-Fluoro-phenylamino)-pyrimidin-4-yl]-4-methyl-thiazol-2-yl}-3-morpholin-4-yl-propionamide
22		N-{4-Methyl-5-[2-(3-nitro-phenylamino)-pyrimidin-4-yl]-thiazol-2-yl}-3-morpholin-4-yl-propionamide
23		N-{4-Methyl-5-[2-(3-nitro-phenylamino)-pyrimidin-4-yl]-thiazol-2-yl}-3-(4-methyl-piperazin-1-yl)-propionamide
24		2-Chloro-N-{4-methyl-5-[2-(3-nitro-phenylamino)-pyrimidin-4-yl]-thiazol-2-yl}-acetamide





25		2-Chloro-N-{5-[2-(3-chloro-phenylamino)-pyrimidin-4-yl]-4-methyl-thiazol-2-yl}-acetamide
26		2-Chloro-N-{5-[2-(3-methoxy-phenylamino)-pyrimidin-4-yl]-4-methyl-thiazol-2-yl}-acetamide
27		2-Chloro-N-{5-[2-(4-dimethylamino-phenylamino)-pyrimidin-4-yl]-4-methyl-thiazol-2-yl}-acetamide
28		4-({4-Methyl-5-[2-(3-nitro-phenylamino)-pyrimidin-4-yl]-thiazol-2-ylcarbamoyl}-methyl)-piperazine-1-carboxylic acid tert-butyl ester







86






29		N-{5-[2-(4-Dimethylamino-phenylamino)-pyrimidin-4-yl]-4-methyl-thiazol-2-yl}-2-[2-(2-hydroxy-ethoxy)-ethylamino]-acetamide
30		6-[5-(2-Oxo-hexahydro-thieno[3,4-d]imidazol-4-yl)-pentanoylamino]-hexanoic acid (2-{4-methyl-5-[2-(4-morpholin-4-yl-phenylamino)-pyrimidin-4-yl]-thiazol-2-ylcarbamoyl}-ethyl)-amide
31		N-{5-[2-(4-Fluoro-phenylamino)-pyrimidin-4-yl]-4-methyl-thiazol-2-yl}-methanesulfonamide
32		N-{4-Methyl-5-[2-(3-nitro-phenylamino)-pyrimidin-4-yl]-thiazol-2-yl}-methane sulfonamide






33		2-Chloro-N-{5-[2-(4-fluoro-phenylamino)-pyrimidin-4-yl]-4-methyl-thiazol-2-yl}-acetamide
34		2-Chloro-N-{5-[2-(4-chloro-phenylamino)-pyrimidin-4-yl]-4-methyl-thiazol-2-yl}-acetamide
35		N-{5-[2-(4-Chloro-phenylamino)-pyrimidin-4-yl]-4-methyl-thiazol-2-yl}-3-morpholin-4-yl-propionamide
36		N-{5-[2-(4-Chloro-phenylamino)-pyrimidin-4-yl]-4-methyl-thiazol-2-yl}-3-(2-diethylamino-ethylamino)-propionamide






88

37		N-{5-[2-(4-Chloro-phenylamino)-pyrimidin-4-yl]-4-methyl-thiazol-2-yl}-3-(2-morpholin-4-yl-ethylamino)-propionamide
38		N-{5-[2-(4-Methoxy-phenylamino)-pyrimidin-4-yl]-4-methyl-thiazol-2-yl}-3-morpholin-4-yl-propionamide
39		N-{5-[2-(3-Methoxy-phenylamino)-pyrimidin-4-yl]-4-methyl-thiazol-2-yl}-3-morpholin-4-yl-propionamide
40		N-{5-[2-(4-Methoxy-phenylamino)-pyrimidin-4-yl]-4-methyl-thiazol-2-yl}-3-(4-methyl-piperazin-1-yl)-propionamide

45		N-{4-[4-(2,4-Dimethyl-thiazol-5-yl)-pyrimidin-2-ylamino]-phenyl}-acetamide
48		N-{4-[4-(2,4-Dimethyl-thiazol-5-yl)-pyrimidin-2-ylamino]-phenyl}-N-methyl-acetamide
49		1-(4-{4-[4-(2,4-Dimethyl-thiazol-5-yl)-pyrimidin-2-ylamino]-phenyl}-piperazin-1-yl)-propan-2-ol
50		2-Chloro-N-{4-[4-(2,4-dimethyl-thiazol-5-yl)-pyrimidin-2-ylamino]-phenyl}-acetamide
51		N-{4-[4-(2,4-Dimethyl-thiazol-5-yl)-pyrimidin-2-ylamino]-phenyl}-2-morpholin-4-yl-acetamide
52		N-{4-[4-(2,4-Dimethyl-thiazol-5-yl)-pyrimidin-2-ylamino]-phenyl}-2-[1,2,4]triazol-1-yl-acetamide

53		N-{4-[4-(2,4-Dimethyl-thiazol-5-yl)-pyrimidin-2-ylamino]-phenyl}-2-pyrrolidin-1-yl-acetamide
54		N-{4-[4-(2,4-Dimethyl-thiazol-5-yl)-pyrimidin-2-ylamino]-phenyl}-2-imidazol-1-yl-acetamide
55		3-[4-(4-Methyl-2-morpholin-4-yl-thiazol-5-yl)-pyrimidin-2-ylamino]-phenol
56		[4-(4-Methyl-2-morpholin-4-yl-thiazol-5-yl)-pyrimidin-2-yl]-(4-morpholin-4-yl-phenyl)-amine
57		N,N-Dimethyl-N'-[4-(4-methyl-2-morpholin-4-yl-thiazol-5-yl)-pyrimidin-2-yl]-benzene-1,4-diamine

59		2-{4-[4-(4-Methyl-2-methylamino-thiazol-5-yl)-pyrimidin-2-ylamino]-phenyl}-ethanol
61		1-(4-{4-[4-(4-Methyl-2-methylamino-thiazol-5-yl)-pyrimidin-2-ylamino]-phenyl}-piperazin-1-yl)-ethanone
62		[4-(4-Methyl-2-methylamino-thiazol-5-yl)-pyrimidin-2-yl]-(4-piperazin-1-yl)-phenyl-amine
63		N-{3-[4-(4-Methyl-2-methylamino-thiazol-5-yl)-pyrimidin-2-ylamino]-benzyl}-acetamide
64		N-{3-[4-(2-Ethylamino-4-methyl-thiazol-5-yl)-pyrimidin-2-ylamino]-benzyl}-acetamide

65		(3-Aminomethyl-phenyl)-[4-(2-ethylamino-4-methyl-thiazol-5-yl)-pyrimidin-2-yl]-amine
66		N-{3-[4-(2,4-Dimethyl-thiazol-5-yl)-pyrimidin-2-ylamino]-benzyl}-acetamide
67		(3-Aminomethyl-phenyl)-[4-(4-methyl-2-methylamino-thiazol-5-yl)-pyrimidin-2-yl]-amine
68		{3-[4-(2-Ethylamino-4-methyl-thiazol-5-yl)-pyrimidin-2-ylamino]-phenyl}-methanol
71		[4-(4-Methyl-2-morpholin-4-yl-thiazol-5-yl)-pyrimidin-2-yl]-(3-nitro-phenyl)-amine

72		{2-Chloro-5-[4-(2,4-dimethyl-thiazol-5-yl)-pyrimidin-2-ylamino]-phenyl}-methanol
73		{2-Chloro-5-[4-(4-methyl-2-methylamino-thiazol-5-yl)-pyrimidin-2-ylamino]-phenyl}-methanol
76		[4-(2-Amino-4-methyl-thiazol-5-yl)-pyrimidin-2-yl]-(4-methanesulfonyl-phenyl)-amine
77		[4-(2-Methoxy-ethoxy)-3-nitro-phenyl]-[4-(4-methyl-2-methylamino-thiazol-5-yl)-pyrimidin-2-yl]-amine
81		[4-(2,4-Dimethyl-thiazol-5-yl)-pyrimidin-2-yl]-[3-(2-morpholin-4-yl-ethoxymethyl)-phenyl]-amine
82		C,C,C-Trifluoro-N-{3-[4-(4-methyl-2-methylamino-thiazol-5-yl)-pyrimidin-2-ylamino]-benzyl}-methanesulfonamide







83		N-{3-[4-(4-Methyl-2-methylamino-thiazol-5-yl)-pyrimidin-2-ylamino]-benzyl}-methanesulfonamide
84		[4-(2-Amino-4-methyl-thiazol-5-yl)-pyrimidin-2-yl]-(3-methanesulfonyl-phenyl)-amine
85		[4-(2,4-Dimethyl-thiazol-5-yl)-pyrimidin-2-yl]-(4-methanesulfonyl-phenyl)-amine
91		{3-[4-(4-Methyl-2-methylamino-thiazol-5-yl)-pyrimidin-2-ylamino]-phenyl}-methanol
92		3,4-Dimethyl-5-[2-(4-morpholin-4-yl-phenylamino)-pyrimidin-4-yl]-3H-thiazol-2-one
93		3,4-Dimethyl-5-{2-[4-(4-methyl-piperazin-1-yl)-phenylamino]-pyrimidin-4-yl}-3H-thiazol-2-one

Table 2. Inhibition of protein kinases by example compounds (refer Table 1).

No.	Kinase Inhibition IC ₅₀ (μM)								
	CDK1 cyclin B ¹	CDK2 cyclin A ¹	CDK2 cyclin E ¹	CDK4 cyclin D1 ¹	CDK7 cyclin H ¹	CDK9 cyclin T1 ¹	GSK- 3β ¹	PLK-1 ¹	ARK- 2 ²
1	17	0.70	0.81	2.4	30	2.7	> 100	98	0.011
2	75	2.2	3.9	3.9	97	23	> 100	> 100	
3	71	4.7	0.43	1.4	106	12	> 100	> 150	
4	30	1.1	0.90	0.43	12	1.8	> 100	> 100	
5	5.6	1.1	0.87	1.7	5.6	6.1	21	35	
6	3.0	0.85	0.71	0.39	1.3	1.0	5.2	13	
7	4.2	1.3	1.8	0.62	2.2				
8	4.9	1.1	1.2	0.36	0.97	0.51	39	> 150	
9	5.2	1.2	2.2	0.43	1.3	0.21			
10	51	4.4	15	11					
11	2.4	2.2	0.26	0.0098	0.019	1.1	6.0	19	
12	4.2	1.0	1.8	0.19	1.1	0.28	53	55	
13	5.6	0.57	0.28	0.67	0.37	0.042			
14			0.068	2.2		0.34			
15	5.4	0.85	0.13	2.0	0.34	0.070	0.61	> 150	
16	2.8	0.65	0.32	6.7	1.1	0.11	0.68	> 150	2.8
17	2.0		0.21						
18			0.18	1.1		0.53			
20	1.9	0.52	0.076	0.77		0.037			
21	0.54	0.32	0.097	1.3	4.9	0.46	14	> 200	
22	0.24	0.098	0.0025	1.7	0.20	0.11	0.25	> 100	
23	3.1	1.4	0.19	3.0	0.81	0.40	1.2	24	
24			1.1	0.44		0.0058			
25	9.4	3.0	0.22	0.97	3.2	0.68	0.083		
28	36	12	2.6	29					
29	0.86	0.49	0.47	1.1	1.2		0.56		
31			0.30	5.6					
32			0.005	1.1		0.0014	0.21		
33			0.23	0.68					
34			1.4	7.0			0.73		
35	15	3.0	0.80	4.7		1.8			
36	33	13	3.4	8.8	4.0	4.8			
37	> 100	16	8.0	18		6.3			
38	1.9	0.58	2.7	1.8		1.1			
39	11	2.7	1.2	32		15			
40	61	14	64	7.4		8.9			
45	2.5	0.84	0.66	0.37	17	0.24			

96

48	12	1.7	2.1	2.6	3.5				
50	3.4	0.89	0.39	0.95	13	0.0018			
51	11	2.9	1.7	0.40	9.6				
52	2.5	1.3	0.44	0.18	2.0				
53	5.2	5.3	0.52	0.040	0.44	0.093	23		
54	1.3	57	0.098	0.032	2.1	0.11	19		
55	2.4	61	0.079	1.0	2.2	2.1	67		
56	18	68	0.60	0.022	0.85	17	> 100		
57	7.7	73	0.35	0.071	0.19	12	74		
59	0.71	0.11	0.14	0.086	2.6	0.47	2.2		
61	2.5	2.8	0.60	0.042	2.3	1.5	6.8		
62	0.98	1.5	0.21	0.0070	0.041	0.098	4.7		
63	48	12	1.7	7.7	20	2.7	8.8	46	
64	0.74	0.31	0.080	0.13	0.59	0.10	2.1	18	
65			0.33	0.11				> 100	
66	0.12	0.10	0.11	0.63	2.1				
67			0.42	0.24				> 100	
68	0.15	0.061	0.065	0.042	0.48		0.82		
71	0.33	0.0060	0.22		2.4	4.2	2.4		
72	62	12	0.73	25	9.6	4.1	48		
73	1.7	1.3	1.0	0.23	2.0	0.55	5.2		
76	1.8	0.14	0.15	3.2	11		5.4		
77	72	1.0	0.15	3.4	1.0	0.26	6.1		
82	2.7	1.3	0.014	2.3	4.0		1.2		
83	0.48	0.15	0.025	0.67	0.37		1.0		
84	0.14	0.05	0.076	0.3	0.43		0.30		
85	0.048	0.001	0.028	3.8	11				
91	0.12	0.10	0.11	0.63					
92	15	1.2	0.64	1.3	2.8		0.47		
93	4.4	1.1	0.84	0.51	0.36	0.3	0.43		

¹, Refer to Table 3 for explanation of abbreviations; ², ARK-2: aurora kinase-2 (also known as aurora A kinase).

Table 3. Kinase specificity of selected compounds (IC₅₀, μ M)

Kinase	Compound				
	1	3	11	15	16
CDK2/E ¹	0.48	0.68	0.26	0.15	0.35
CDK2/A ²	0.44	0.44	2.2	0.40	0.65
CDK1/B1 ³	21	>100	2.4	1.6	2.8
CDK4/D1 ⁴	2.2	0.15	0.0098	1.6	6.7
CDK7/H ⁵	56	>100	0.019	0.39	0.97
CDK9/T1 ⁶	2.3	8.5	1.1	0.11	0.20
ERK2 ⁷	>100	>100	>100	>100	>100
p70/S6 ⁸	2.3	2.6	>100	>100	8.0
CK2 ⁹	>100	>100	>100	>100	>100
PKC α ¹⁰	>100	>100	>100	>100	53
Akt/PKB ¹¹	>100	>100	>100	>100	>100
PKA ¹²	5.8	39	11	2.0	4.9
SAPK2a ¹³	>100	>100	>100	>100	>100
PLK1 ¹⁴	>100	>100	19	>100	>100
CaMKII ¹⁵	26	>100	56	3.9	11
Abl ¹⁶	>100	50	72	0.76	1.5
GSK-3 ¹⁷	>100	>100	6.0	0.54	0.68

¹, CDK2 / cyclin E complex; ², CDK2 / cyclin A complex; ³, CDK 1 / cyclin B1 complex; ⁴, CDK4 / cyclin D1 complex; ⁵, CDK7 / cyclin H / MAT 1 complex; ⁶, CDK9 / cyclin T1 complex; ⁷, extracellular-signal-regulated kinase 2; ⁸, p70 ribosomal protein S6 kinase; ⁹, casein kinase 2; ¹⁰, protein kinase C α ; ¹¹, protein kinase B; ¹², cAMP-dependent protein kinase; ¹³, stress-activated protein kinase 2a; ¹⁴, polo-like kinase 1; ¹⁵, calmodulin-dependent kinase II; ¹⁶, Ablason tyrosine kinase; ¹⁷, glycogen synthase kinase 3 β .

Table 4. Anti-proliferative activity against human cancer cell lines (refer Table 1)

No.	72-h MTT IC ₅₀ (μM)		
	A549	HT29	Saos-2
1	2.1	1.7	1.9
2	3.5	3.3	4.8
3	3.7	2.8	3.1
4	0.77	0.92	1.2
5	3.8	2.2	3.9
6	1.3	1.1	0.78
7	3.9	1.6	1.6
8	0.61	0.80	0.38
9	3.0	2.0	2.0
10	11	6.4	
11	2.2	1.6	3.6
12	0.71	0.74	0.43
13	3.1	2.8	1.9
14	0.99	0.73	1.5
15	1.1	1.0	1.9
16	0.53	0.29	
17	1.1	1.2	1.2
18	2.5	2.4	1.1
20	5.8	6.3	
21	8.6	4.1	4.3
22	0.81	0.52	0.60
23	3.8	1.1	4.4
24	0.14	0.14	0.17
25	0.97	1.3	1.6
28	9.9	6.74	15
29	0.69	2.6	0.72
31	2.0	4.1	0.81
32	0.10	0.17	0.16
33	0.33	0.11	0.30
34	3.9	3.2	3.3
35	15	7.7	26
36	26	9.6	38
37	35	15	73
38	2.7	1.7	5.3
39	0.80	0.72	0.95
40	4.4	0.73	4.3
45	2.6	2.2	2.8
47	2.7	0.35	1.6

48	3.0	2.6	3.8
50	0.16	0.16	0.16
51	1.9	1.4	1.1
52	1.6	1.6	1.1
53	0.77	0.95	0.53
54	2.9	2.7	1.9
55	2.3	5.1	1.4
56	4.4	2.9	5.0
57	2.7	0.24	3.8
59	2.4	1.7	1.9
61	0.72	0.53	1.0
62	0.19	0.18	0.26
63	0.98	1.4	1.2
64	0.19	0.41	0.34
65	3.2	4.8	0.65
67	14	14	1.1
71	80	71	10
72	32	20	18
73	1.5	5.5	2.9
76	5.8	1.8	1.3
77	0.31	0.21	0.35

Table 5. *In vitro* antiproliferative activity of selected compounds (72-h MTT, IC₅₀, μ M)

Cell line		Compound			
Type	Designation	1	11	15	16
Bone osteosarcoma	Saos-2	0.1	3.7	0.4	0.6
Bone osteosarcoma	U2OS	2.1	2.3	1.02	0.48
Breast	MCF-7	> 5	1.9	0.9	0.39
Cervix	Hela	1.8	6.2	0.7	0.4
Colon	HT29	1.1	1.6	0.5	0.3
Colon	Lovo	0.9	2.0	0.7	0.3
Colon	H1299	0.9	1.1	1.3	0.6
Colon	HCT-116	0.9	0.6	0.6	0.3
Gastric adenocarcinoma	AGS	1.1	1.3	1.1	0.3
Leiomyosarcoma	SKUT-1B		0.3	0.2	0.1
Leiomyosarcoma	SKUT-1	0.9	0.8	0.9	0.2
Chronic myelogenous leukaemia	K562	3.8	2.1	4.6	2.5
Leukemia	CCRF-CEM	0.9	0.5	2.6	1.1
Promyelocytic leukaemia	HL60	1.8	1.7	2.2	0.6
Lung	nci-H460	0.2	0.7	1.2	0.3
Lung	A549	1.0	2.2	0.6	0.5
Neuroblastoma	SK-N-MC	0.3	0.6	0.5	0.4
Osteogenic sarcoma	SJSA-1	> 5	4.3	1.9	1.0
Prostate	DU-145	1.5	1.0	1.3	0.6
Skin keratinocytes	Hacat	1.1	1.0	1.5	1.1
Uterine	Messa	0.2	0.1	0.9	0.4
Uterine	Messa-Dx5	0.2	0.2	0.4	0.1
<i>Average (all transformed cells)</i>		<i>1.1</i>	<i>1.6</i>	<i>1.2</i>	<i>0.6</i>
<i>SD (all transformed cells)</i>		<i>0.9</i>	<i>1.4</i>	<i>1.0</i>	<i>0.5</i>
<i>Median (all transformed cells)</i>		<i>1.0</i>	<i>1.1</i>	<i>0.9</i>	<i>0.4</i>
Foreskin fibroblast (non-transformed)	Hs27	> 5	19	1.7	> 5
Foetal lung fibroblast (non-transformed)	IMR-90	> 5	31	2.5	1.7
Foetal lung fibroblast (non-transformed)	WI38	> 5	22	> 5	1.4


Table 6. Summary of anti-HIV activity

<u>Compound</u>	HIV-1 / PBMC		PBMC TC₅₀ (μM)	TI
	IC₅₀ (nM)	IC₉₀ (nM)		
AZT ^a	4	10	> 1	> 231
1	1,327	2,645	6.1	4.6
3	297	679	> 100	> 337
4	166	295	9.4	57
15	124	385	2.4	20
53	812	871	1.6	1.9

^a, AZT: Azidothymidine; anti-HIV drug in clinical use as positive control.

102
CLAIMS

1. A compound of formula I, or a pharmaceutically acceptable salt thereof,



wherein:

- (A) one of X^1 and X^2 is S, and the other of X^1 and X^2 is N;
 "a" is a single bond; and
 "b", "c", "d", "e" and "f" are single or double bonds so as to form a thiazolyl ring;
 R^2 is independently as defined below for R^1 , R^{3-8} ; or
- (B) one of X^1 and X^2 is S, and the other of X^1 and X^2 is NR^{17} ;
 "a" and "d" are each double bonds; and
 "b", "c", "e" and "f" are each single bonds;
 R^2 is oxo;
 R^{17} is H or alkyl;

where:

Z is NH, NHCO, NHSO₂, NHCH₂, CH₂, CH₂CH₂, CH=CH, SO₂, or SO;

R^1 , R^3 , R^4 , R^5 , R^6 , R^7 and R^8 are each independently H, alkyl, alkyl- R^9 , aryl, aryl- R^9 , aralkyl, aralkyl- R^9 , halogeno, NO₂, CN, OH, O-alkyl, COR⁹, COOR⁹, O-aryl, O-R⁹, NH₂, NH-alkyl, NH-aryl, N-(alkyl)₂, N-(aryl)₂, N-(alkyl)(aryl), NH-R⁹, N-(R⁹)(R¹⁰), N-

103

(alkyl)(R⁹), N-(aryl)(R⁹), COOH, CONH₂, CONH-alkyl, CONH-aryl, CON-(alkyl)(R⁹), CON(aryl)(R⁹), CONH-R⁹, CON-(R⁹)(R¹⁰), SO₃H, SO₂-alkyl, SO₂-alkyl-R⁹, SO₂-aryl, SO₂-aryl-R⁹, SO₂NH₂, SO₂NH-R⁹, SO₂N-(R⁹)(R¹⁰), CF₃, CO-alkyl, CO-alkyl-R⁹, CO-aryl, CO-aryl-R⁹ or R¹¹, wherein alkyl, aryl, aralkyl groups may be further substituted with one or more groups selected from halogeno, NO₂, OH, O-methyl, NH₂, COOH, CONH₂ and CF₃;

wherein at least one of R¹, R², R³, R⁴, R⁵, R⁶, R⁷ and R⁸ is an R⁹ or R¹⁰-containing group, or is R¹¹;

R⁹ and R¹⁰ are each independently solubilising groups selected from:

- (i)
 - a mono-, di- or polyhydroxylated alicyclic group;
 - a di- or polyhydroxylated aliphatic or aromatic group;
 - a carbohydrate derivative;
 - an O- and/or S-containing heterocyclic group optionally substituted by one or more hydroxyl groups;
 - an aliphatic or aromatic group containing a carboxamide, sulfoxide, sulfone, or sulfonamide function; or
 - a halogenated alkylcarbonyl group;
- (ii) COOH, SO₃H, OSO₃H, PO₃H₂, or OPO₃H₂;
- (iii) Y, where Y is selected from an alicyclic, aromatic, or heterocyclic group comprising one or more of the functions =N-, -O-, -NH₂, -NH-, a quarternary amine salt, guanidine, and amidine, where Y is optionally substituted by one or more substituents selected from:
 - SO₂-alkyl;
 - alkyl optionally substituted by one or more OH groups;
 - CO-alkyl;
 - aralkyl;
 - COO-alkyl; and
 - an ether group optionally substituted by one or more OH groups; and

where Y is other than pyridinyl;

- (iv) a natural or unnatural amino acid, a peptide or a peptide derivative;

each R^{11} is a solubilising group as defined for R^9 and R^{10} in (i) or (iv) above; or is selected from:

- (v) OSO_3H , PO_3H_2 , or OPO_3H_2 ;
- (vi) Y as defined above, but excluding guanidine and quarternary amine salts;
- (vii) $NHCO(CH_2)_m[NHCO(CH_2)_{m'}]_p[NHCO(CH_2)_{m''}]_qY$ or $NHCO(CH_2)_tNH(CH_2)_{t'}Y$ where p and q are each 0 or 1, and m, m', m'', t and t' are each independently an integer from 1 to 10; and
- (viii) $(CH_2)_nNR^{14}COR^{12}$, $(CH_2)_nNR^{15}SO_2R^{13}$, or SO_2R^{16} , where R^{12} , R^{13} and R^{16} are each alkyl groups optionally comprising one or more heteroatoms, and which are optionally substituted by one or more substituents selected from OH, NH_2 , halogen and NO_2 , R^{14} and R^{15} are each independently H or alkyl, and n and n' are each independently 0, 1, 2, or 3;
- (ix) an ether or polyether optionally substituted by one or more hydroxyl groups or one or more Y groups;
- (x) $(CH_2)_rNH_2$; where r is 0, 1, 2, or 3;
- (xi) $(CH_2)_{r'}OH$; where r' is 0, 1, 2, or 3;

with the proviso that the compound is other than:

2-Chloro-N-{4-methyl-5-[2-(3-nitro-phenylamino)-pyrimidin-4-yl]-thiazol-2-yl}-acetamide;

N-{4-Methyl-5-[2-(3-nitro-phenylamino)-pyrimidin-4-yl]-thiazol-2-yl}-methane sulfonamide;


2-Chloro-N-{5-[2-(4-fluoro-phenylamino)-pyrimidin-4-yl]-4-methyl-thiazol-2-yl}-acetamide;

{3-[4-(2,4-Dimethyl-thiazol-5-yl)-pyrimidin-2-ylamino]-phenyl}-methanol;

2-{4-[4-(2-Amino-4-methyl-thiazol-5-yl)-pyrimidin-2-ylamino]-phenyl}-ethanol; or

2-{4-[4-(2,4-Dimethyl-thiazol-5-yl)-pyrimidin-2-ylamino]-phenyl}-ethanol.

2. A compound according to claim 1 of formula Ib, or a pharmaceutically acceptable salt thereof,



Ib

wherein one of X¹ and X² is S, and the other of X¹ and X² is N, and Z, R¹⁻⁸ are as defined in claim 1.

3. A compound according to claim 2 wherein X¹ is S and X² is N.
4. A compound according to any preceding claim wherein Z is NH.
5. A compound according to any preceding claim wherein R³ is H.
6. A compound according to any preceding claim wherein at least one of R², R⁵, R⁶ or R⁷ is an R⁹ or R¹⁰-containing group, or is R¹¹.
7. A compound according to any preceding claim wherein X¹ is S, X² is N, Z is NH, R¹ is Me, R² is alkyl or amino, R³ is H, one or two of R⁵, R⁶, and R⁷ are CF₃, OH, O-alkyl, halogeno, NO₂, NH₂, NH-alkyl or N-(alkyl)₂ and at least one of R², R⁵, R⁶ or R⁷ is an R⁹ or R¹⁰-containing group, or is R¹¹.
8. A compound according to any one of claims 1 to 5 wherein at least one of R¹, R², R³, R⁴, R⁵, R⁶, R⁷ and R⁸ is R¹¹.

9. A compound according to any preceding claim wherein the solubilising group is R^{11} , and R^{11} is:

- (b) Y as defined in claim 1, but excluding guanidine, where Y can also be an alicyclic, aromatic, or heterocyclic group comprising one or more =N- groups;
- (b) $NHCO(CH_2)_m[NHCO(CH_2)_{m'}]_p[NHCO(CH_2)_{m''}]_qY$ or $NHCO(CH_2)_tNH(CH_2)_{t'}Y$ where p and q are each 0 or 1, and m, m', m'', t and t' are each an integer from 1 to 10; or
- (c) $(CH_2)_nNR^{14}COR^{12}$, $(CH_2)_nNR^{15}SO_2R^{13}$, or SO_2R^{16} , where R^{12} , R^{13} and R^{16} are each alkyl groups optionally comprising one or more heteroatoms, and which are substituted by one or more substituents selected from OH, NH_2 , halogen and NO_2 , R^{14} and R^{15} are each independently H or alkyl, and n and n' are each independently 0, 1, 2, or 3.

10. A compound according to claim 9 where Y is an alicyclic group comprising one or more of the functions -O-, NH_2 , -NH-, =N-, a quarternary amine salt, or amidine, and wherein Y is optionally substituted by one or more substituents as defined in claim 1.


11. A compound according to any preceding claim wherein Y is a morpholine or piperazine group, each of which may be optionally substituted by one or more substituents selected from SO_2 -alkyl, alkyl optionally substituted by one or more OH groups, CO-alkyl, aralkyl, COO-alkyl, and an ether group optionally substituted by one or more OH groups


12. A compound according to any one of claims 1 to 9 wherein Y is a 2-oxo-hexahydro-thien[3,4-d]imidazole group.

13. A compound according to any preceding claim wherein at least one of R^2 , R^6 or R^7 is R^{11} .

107


14. A compound according to any preceding claim wherein R¹¹ is selected from the following:






15. A compound according to claim 13 or claim 14 wherein R^6 or R^7 is R^{11} .
16. A compound according to claim 15 wherein R^6 is R^{11} and R^2 , R^4 , R^5 , R^7 and R^8 are each independently selected from alkyl, H, CF_3 , OH, O-alkyl, halogeno, NO_2 , NH_2 , NH-alkyl and N-(alkyl)₂.
17. A compound according to claim 15 or claim 16 wherein R^6 is R^{11} and R^2 , R^4 , R^5 , R^7 and R^8 are each independently selected from alkyl, H, O-alkyl, halogeno, NO_2 , NH_2 and NH-alkyl.
18. A compound according to any one of claims 15 to 17 wherein R^6 is R^{11} and R^4 , R^5 , R^7 and R^8 are all H and R^2 is selected from alkyl, O-alkyl, NH_2 and NH-alkyl.
19. A compound according to any one of claims 15 to 18 wherein R^{11} is selected from:

109



20. A compound according to claim 15 wherein R⁷ is R¹¹ and R⁴, R⁵, R⁶, R⁸ are all H, and R² is selected from alkyl, O-alkyl, NH₂ and NH-alkyl.

21. A compound according to claim 20 wherein R¹¹ is selected from:



22. A compound according to claim 13 or claim 14 wherein R² is R¹¹.


23. A compound according to claim 22 wherein R² is R¹¹ and R⁴, R⁵, R⁶, R⁷ and R⁸ are each independently selected from alkyl, H, CF₃, OH, O-alkyl, halogeno, NO₂, NH₂, NH-alkyl and N-(alkyl)₂.

24. A compound according to claim 22 or claim 23 wherein R² is R¹¹ and R⁴, R⁵, R⁶, R⁷ and R⁸ are each independently selected from H, O-alkyl, halogeno, N-(alkyl)₂, NO₂.

110

25. A compound according to any one of claims 22 to 24 wherein R^2 is R^{11} , one of R^5 or R^7 is selected from NO_2 , alkoxy, halogeno and $\text{N}(\text{alkyl})_2$, and the remainder of R^4 , R^5 , R^6 , R^7 and R^8 are all H.


26. A compound according to any one of claims 22 to 24 wherein R^{11} is selected from:



27. A compound according to any preceding claim wherein R^1 is methyl, Z is NH and R^3 is H.

111

28. A compound according to claim 1 of formula Id, or or a pharmaceutically acceptable salt thereof,



Id

29. A compound according to claim 28 wherein R¹ and R³⁻⁸ are as defined in any one of claims 4, 9-12, 14-17, 19, 21

30. A compound according to any preceding claim which is selected from the following:


1	[4-(2-Amino-4-methyl-thiazol-5-yl)-pyrimidin-2-yl]-(4-morpholin-4-yl-phenyl)-amine
2	[4-(2,4-Dimethyl-thiazol-5-yl)-pyrimidin-2-yl]-(4-morpholin-4-yl-phenyl)-amine
3	[4-(2-N-Methylamino-4-methyl-thiazol-5-yl)-pyrimidin-2-yl]-(4-morpholinophenyl)-amine
4	[4-(2-Ethylamino-4-methyl-thiazol-5-yl)-pyrimidin-2-yl]-(4-morpholin-4-yl-phenyl)-amine
5	1-(4-{4-[4-(2,4-Dimethyl-thiazol-5-yl)-pyrimidin-2-ylamino]-phenyl}-piperazin-1-yl)-ethanone
6	[4-(2,4-Dimethyl-thiazol-5-yl)-pyrimidin-2-yl]-(4-piperazin-1-yl-phenyl)-amine
7	[4-(2,4-Dimethyl-thiazol-5-yl)-pyrimidin-2-yl]-[4-(4'-2''-ethoxyethanolpiperazino)-phenyl]-amine
8	3-(4-{4-[4-(2,4-Dimethyl-thiazol-5-yl)-pyrimidin-2-ylamino]-phenyl}-piperazin-1-yl)-propan-1-ol
9	2-(4-{4-[4-(2,4-Dimethyl-thiazol-5-yl)-pyrimidin-2-ylamino]-phenyl}-piperazin-1-yl)-ethanol
10	[4-(2,4-Dimethyl-thiazol-5-yl)-pyrimidin-2-yl]-[4-(4-methanesulfonyl-piperazin-1-yl)-phenyl]-amine
11	[4-(4-Benzyl-piperazin-1-yl)-phenyl]-[4-(4-methyl-2-methylamino-thiazol-5-yl)-pyrimidin-2-yl]-amine
12	[4-(2,4-Dimethyl-thiazol-5-yl)-pyrimidin-2-yl]-[4-(4-methyl-piperazin-1-yl)-phenyl]-amine

13	3-Amino-N-{4-methyl-5-[2-(3-nitro-phenylamino)-pyrimidin-4-yl]-thiazol-2-yl}-propionamide
14	(2S)-2-Amino-3-hydroxy-N-{4-methyl-5-[2-(3-nitro-phenylamino)-pyrimidin-4-yl]-thiazol-2-yl}-propionamide
15	(2R,3R)-2-Amino-3-hydroxy-N-{4-methyl-5-[2-(3-nitro-phenylamino)-pyrimidin-4-yl]-thiazol-2-yl}-butyramide
16	(2R)-2-Amino-N-{4-methyl-5-[2-(3-nitro-phenylamino)-pyrimidin-4-yl]-thiazol-2-yl}-butyramide
17	(2S,3S)-2-Amino-3-hydroxy-N-{4-methyl-5-[2-(3-nitro-phenylamino)-pyrimidin-4-yl]-thiazol-2-yl}-butyramide
18	4-Amino-N-{4-methyl-5-[2-(3-nitro-phenylamino)-pyrimidin-4-yl]-thiazol-2-yl}-butyramide
19	3-Amino-N-{4-methyl-5-[2-(4-morpholin-4-yl-phenylamino)-pyrimidin-4-yl]-thiazol-2-yl}-propionamide
20	3-Bromo-N-{5-[2-(4-fluoro-phenylamino)-pyrimidin-4-yl]-4-methyl-thiazol-2-yl}-propionamide
21	N-{5-[2-(4-Fluoro-phenylamino)-pyrimidin-4-yl]-4-methyl-thiazol-2-yl}-3-morpholin-4-yl-propionamide
22	N-{4-Methyl-5-[2-(3-nitro-phenylamino)-pyrimidin-4-yl]-thiazol-2-yl}-3-morpholin-4-yl-propionamide
23	N-{4-Methyl-5-[2-(3-nitro-phenylamino)-pyrimidin-4-yl]-thiazol-2-yl}-3-(4-methyl-piperazin-1-yl)-propionamide
24	2-Chloro-N-{4-methyl-5-[2-(3-nitro-phenylamino)-pyrimidin-4-yl]-thiazol-2-yl}-acetamide
25	2-Chloro-N-{5-[2-(3-chloro-phenylamino)-pyrimidin-4-yl]-4-methyl-thiazol-2-yl}-acetamide
26	2-Chloro-N-{5-[2-(3-methoxy-phenylamino)-pyrimidin-4-yl]-4-methyl-thiazol-2-yl}-acetamide
27	2-Chloro-N-{5-[2-(4-dimethylamino-phenylamino)-pyrimidin-4-yl]-4-methyl-thiazol-2-yl}-acetamide
28	4-({4-Methyl-5-[2-(3-nitro-phenylamino)-pyrimidin-4-yl]-thiazol-2-ylcarbamoyl}-methyl)-piperazine-1-carboxylic acid tert-butyl ester
29	N-{5-[2-(4-Dimethylamino-phenylamino)-pyrimidin-4-yl]-4-methyl-thiazol-2-yl}-2-[2-(2-hydroxy-ethoxy)-ethylamino]-acetamide
30	6-[5-(2-Oxo-hexahydro-thieno[3,4-d]imidazol-4-yl)-pentanoylamino]-hexanoic acid (2-{4-methyl-5-[2-(4-morpholin-4-yl-phenylamino)-pyrimidin-4-yl]-thiazol-2-ylcarbamoyl}-ethyl)-amide
31	N-{5-[2-(4-Fluoro-phenylamino)-pyrimidin-4-yl]-4-methyl-thiazol-2-yl}-methanesulfonamide
32	N-{4-Methyl-5-[2-(3-nitro-phenylamino)-pyrimidin-4-yl]-thiazol-2-yl}-methane sulfonamide
33	2-Chloro-N-{5-[2-(4-fluoro-phenylamino)-pyrimidin-4-yl]-4-methyl-thiazol-2-yl}-acetamide

34	2-Chloro-N-{5-[2-(4-chloro-phenylamino)-pyrimidin-4-yl]-4-methyl-thiazol-2-yl}-acetamide
35	N-{5-[2-(4-Chloro-phenylamino)-pyrimidin-4-yl]-4-methyl-thiazol-2-yl}-3-morpholin-4-yl-propionamide
36	N-{5-[2-(4-Chloro-phenylamino)-pyrimidin-4-yl]-4-methyl-thiazol-2-yl}-3-(2-diethylamino-ethylamino)-propionamide
37	N-{5-[2-(4-Chloro-phenylamino)-pyrimidin-4-yl]-4-methyl-thiazol-2-yl}-3-(2-morpholin-4-yl-ethylamino)-propionamide
38	N-{5-[2-(4-Methoxy-phenylamino)-pyrimidin-4-yl]-4-methyl-thiazol-2-yl}-3-morpholin-4-yl-propionamide
39	N-{5-[2-(3-Methoxy-phenylamino)-pyrimidin-4-yl]-4-methyl-thiazol-2-yl}-3-morpholin-4-yl-propionamide
40	N-{5-[2-(4-Methoxy-phenylamino)-pyrimidin-4-yl]-4-methyl-thiazol-2-yl}-3-(4-methyl-piperazin-1-yl)-propionamide
45	N-{4-[4-(2,4-Dimethyl-thiazol-5-yl)-pyrimidin-2-ylamino]-phenyl}-acetamide
48	N-{4-[4-(2,4-Dimethyl-thiazol-5-yl)-pyrimidin-2-ylamino]-phenyl}-N-methyl-acetamide
49	1-(4-{4-[4-(2,4-Dimethyl-thiazol-5-yl)-pyrimidin-2-ylamino]-phenyl}-piperazin-1-yl)-propan-2-ol
50	2-Chloro-N-{4-[4-(2,4-dimethyl-thiazol-5-yl)-pyrimidin-2-ylamino]-phenyl}-acetamide
51	N-{4-[4-(2,4-Dimethyl-thiazol-5-yl)-pyrimidin-2-ylamino]-phenyl}-2-morpholin-4-yl-acetamide
52	N-{4-[4-(2,4-Dimethyl-thiazol-5-yl)-pyrimidin-2-ylamino]-phenyl}-2-[1,2,4]triazol-1-yl-acetamide
53	N-{4-[4-(2,4-Dimethyl-thiazol-5-yl)-pyrimidin-2-ylamino]-phenyl}-2-pyrrolidin-1-yl-acetamide
54	N-{4-[4-(2,4-Dimethyl-thiazol-5-yl)-pyrimidin-2-ylamino]-phenyl}-2-imidazol-1-yl-acetamide
55	3-[4-(4-Methyl-2-morpholin-4-yl-thiazol-5-yl)-pyrimidin-2-ylamino]-phenol
56	[4-(4-Methyl-2-morpholin-4-yl-thiazol-5-yl)-pyrimidin-2-yl]-(4-morpholin-4-yl-phenyl)-amine
57	N,N-Dimethyl-N'-[4-(4-methyl-2-morpholin-4-yl-thiazol-5-yl)-pyrimidin-2-yl]-benzene-1,4-diamine
59	2-{4-[4-(4-Methyl-2-methylamino-thiazol-5-yl)-pyrimidin-2-ylamino]-phenyl}-ethanol
61	1-(4-{4-[4-(4-Methyl-2-methylamino-thiazol-5-yl)-pyrimidin-2-ylamino]-phenyl}-piperazin-1-yl)-ethanone
62	[4-(4-Methyl-2-methylamino-thiazol-5-yl)-pyrimidin-2-yl]-(4-piperazin-1-yl-phenyl)-amine
63	N-{3-[4-(4-Methyl-2-methylamino-thiazol-5-yl)-pyrimidin-2-ylamino]-benzyl}-acetamide

64	N-{3-[4-(2-Ethylamino-4-methyl-thiazol-5-yl)-pyrimidin-2-ylamino]-benzyl}-acetamide
65	(3-Aminomethyl-phenyl)-[4-(2-ethylamino-4-methyl-thiazol-5-yl)-pyrimidin-2-yl]-amine
66	N-{3-[4-(2,4-Dimethyl-thiazol-5-yl)-pyrimidin-2-ylamino]-benzyl}-acetamide
67	(3-Aminomethyl-phenyl)-[4-(4-methyl-2-methylamino-thiazol-5-yl)-pyrimidin-2-yl]-amine
68	{3-[4-(2-Ethylamino-4-methyl-thiazol-5-yl)-pyrimidin-2-ylamino]-phenyl}-methanol
71	[4-(4-Methyl-2-morpholin-4-yl-thiazol-5-yl)-pyrimidin-2-yl]-(3-nitro-phenyl)-amine
72	{2-Chloro-5-[4-(2,4-dimethyl-thiazol-5-yl)-pyrimidin-2-ylamino]-phenyl}-methanol
73	{2-Chloro-5-[4-(4-methyl-2-methylamino-thiazol-5-yl)-pyrimidin-2-ylamino]-phenyl}-methanol
76	[4-(2-Amino-4-methyl-thiazol-5-yl)-pyrimidin-2-yl]-(4-methanesulfonyl-phenyl)-amine
77	[4-(2-Methoxy-ethoxy)-3-nitro-phenyl]-[4-(4-methyl-2-methylamino-thiazol-5-yl)-pyrimidin-2-yl]-amine
81	[4-(2,4-Dimethyl-thiazol-5-yl)-pyrimidin-2-yl]-[3-(2-morpholin-4-yl-ethoxymethyl)-phenyl]-amine
82	C,C,C-Trifluoro-N-{3-[4-(4-methyl-2-methylamino-thiazol-5-yl)-pyrimidin-2-ylamino]-benzyl}-methanesulfonamide
83	N-{3-[4-(4-Methyl-2-methylamino-thiazol-5-yl)-pyrimidin-2-ylamino]-benzyl}-methanesulfonamide
84	[4-(2-Amino-4-methyl-thiazol-5-yl)-pyrimidin-2-yl]-(3-methanesulfonyl-phenyl)-amine
85	[4-(2,4-Dimethyl-thiazol-5-yl)-pyrimidin-2-yl]-(4-methanesulfonyl-phenyl)-amine
91	{3-[4-(4-Methyl-2-methylamino-thiazol-5-yl)-pyrimidin-2-ylamino]-phenyl}-methanol
92	3,4-Dimethyl-5-[2-(4-morpholin-4-yl-phenylamino)-pyrimidin-4-yl]-3H-thiazol-2-one
93	3,4-Dimethyl-5-{2-[4-(4-methyl-piperazin-1-yl)-phenylamino]-pyrimidin-4-yl}-3H-thiazol-2-one

31. A compound according to any preceding claim which is selected from the following: [4], [8], [12], [14], [16], [22], [24], [25], [29], [32], [33], [39], [40], [50], [53], [61], [57], [62], [63], [64], [65] and [77].
32. A compound according to claim 31 which is selected from the following: [24], [25], [32], [33], [50], [62] and [64].
33. A compound according to any one of claims 1 to 30 which is selected from the following: [11], [13], [14], [15], [20], [21], [22], [24], [25], [32], [50], [53], [54], [55], [56], [57], [59], [61], [62], [64], [68], [71], [82], 83, [84] and [85].
34. A compound according to claim 33 which is selected from the following: [11], [22], [24], [32], [50], [62], [71] and [85].
35. A pharmaceutical composition comprising a compound according to any one of claims 1 to 34 admixed with a pharmaceutically acceptable diluent, excipient or carrier.
36. Use of a compound according to any one of claims 1 to 26 in the preparation of a medicament for treating a proliferative disorder.
37. Use according to claim 36 wherein the proliferative disorder is cancer or leukaemia.
38. Use according claim 36 or claim 37 wherein said compound is administered in combination with one or more other anticancer compounds.
39. Use of a compound of formula Ia, or a pharmaceutically acceptable salt thereof,



wherein:

- (A) one of X^1 and X^2 is S, and the other of X^1 and X^2 is N;
 "a" is a single bond; and
 "b", "c", "d", "e" and "f" are single or double bonds so as to form a thiazolyl ring;
 R^2 is independently as defined below for R^1 , R^{3-8} ; or
- (B) one of X^1 and X^2 is S, and the other of X^1 and X^2 is NR^{17} ;
 "a" and "d" are each double bonds; and
 "b", "c", "e" and "f" are each single bonds;
 R^2 is oxo;
 R^{17} is H or alkyl;

where:

Z is NH, NHCO, NHSO₂, NHCH₂, CH₂, CH₂CH₂, CH=CH, SO₂, or SO;

R^1 , R^3 , R^4 , R^5 , R^6 , R^7 and R^8 are each independently H, alkyl, alkyl- R^9 , aryl, aryl- R^9 , aralkyl, aralkyl- R^9 , halogeno, NO₂, CN, OH, O-alkyl, COR⁹, COOR⁹, O-aryl, O- R^9 , NH₂, NH-alkyl, NH-aryl, N-(alkyl)₂, N-(aryl)₂, N-(alkyl)(aryl), NH- R^9 , N-(R^9)(R^{10}), N-(alkyl)(R^9), N-(aryl)(R^9), COOH, CONH₂, CONH-alkyl, CONH-aryl, CON-(alkyl)(R^9), CON(aryl)(R^9), CONH- R^9 , CON-(R^9)(R^{10}), SO₃H, SO₂-alkyl, SO₂-alkyl- R^9 , SO₂-aryl, SO₂-aryl- R^9 , SO₂NH₂, SO₂NH- R^9 , SO₂N-(R^9)(R^{10}), CF₃, CO-alkyl, CO-alkyl- R^9 , CO-aryl, CO-aryl- R^9 or R^{11} , wherein alkyl, aryl, aralkyl groups may be further substituted with one

117

or more groups selected from halogeno, NO₂, OH, O-methyl, NH₂, COOH, CONH₂ and CF₃;

wherein at least one of R¹, R², R³, R⁴, R⁵, R⁶, R⁷ and R⁸ is an R⁹ or R¹⁰-containing group, or is R¹¹;

R⁹ and R¹⁰ are each independently solubilising groups selected from:

- (i)
 - a mono-, di- or polyhydroxylated alicyclic group;
 - a di- or polyhydroxylated aliphatic or aromatic group;
 - a carbohydrate derivative;
 - an O- and/or S-containing heterocyclic group optionally substituted by one or more hydroxyl groups;
 - an aliphatic or aromatic group containing a carboxamide, sulfoxide, sulfone, or sulfonamide function; or
 - a halogenated alkylcarbonyl group;
- (ii) COOH, SO₃H, OSO₃H, PO₃H₂, or OPO₃H₂;
- (iii) Y, where Y is selected from an alicyclic, aromatic, or heterocyclic group comprising one or more of the functions =N-, -O-, -NH₂, -NH-, a quarternary amine salt, guanidine, and amidine, where Y is optionally substituted by one or more substituents selected from:
 - SO₂-alkyl;
 - alkyl optionally substituted by one or more OH groups;
 - CO-alkyl;
 - aralkyl;
 - COO-alkyl; and
 - an ether group optionally substituted by one or more OH groups; andwhere Y is other than pyridinyl;
- (iv) a natural or unnatural amino acid, a peptide or a peptide derivative;

118

each R^{11} is a solubilising group as defined for R^9 and R^{10} in (i) or (iv) above; or is selected from:

- (v) OSO_3H , PO_3H_2 , or OPO_3H_2 ;
- (vi) Y as defined above, but excluding guanidine and quarternary amine salts;
- (vii) $NHCO(CH_2)_m[NHCO(CH_2)_{m'}]_p[NHCO(CH_2)_{m''}]_qY$ or $NHCO(CH_2)_tNH(CH_2)_{t'}Y$ where p and q are each 0 or 1, and m, m', m'', t and t' are each independently an integer from 1 to 10; and
- (viii) $(CH_2)_nNR^{14}COR^{12}$, $(CH_2)_nNR^{15}SO_2R^{13}$, or SO_2R^{16} , where R^{12} , R^{13} and R^{16} are each alkyl groups optionally comprising one or more heteroatoms, and which are optionally substituted by one or more substituents selected from OH, NH_2 , halogen and NO_2 , R^{14} and R^{15} are each independently H or alkyl, and n and n' are each independently 0, 1, 2, or 3;
- (ix) an ether or polyether optionally substituted by one or more hydroxyl groups or one or more Y groups;
- (x) $(CH_2)_rNH_2$; where r is 0, 1, 2, or 3;
- (xi) $(CH_2)_{r'}OH$; where r' is 0, 1, 2, or 3;

in the preparation of a medicament for treating a viral disorder.

40. Use according to claim 39 wherein said compound of formula Ia is as defined in any one of claims 2 to 30.


41. Use according to claim 39 wherein said compound of formula Ia is as defined in claim 30 or is additionally selected from the following:

N-{4-Methyl-5-[2-(3-nitro-phenylamino)-pyrimidin-4-yl]-thiazol-2-yl}
methanesulfonamide [32];

2-Chloro-N-{4-methyl-5-[2-(3-nitro-phenylamino)-pyrimidin-4-yl]-thiazol-2-yl}-
acetamide [24]; and

2-Chloro-N-{5-[2-(4-fluoro-phenylamino)-pyrimidin-4-yl]-4-methyl-thiazol-2-yl}-
acetamide [33].

42. Use according to any one of claims 39 to 41 wherein said compound of formula Ia is selected from the following: [1], [3], [4], [15] and [53].
43. Use according to any one of claims 39 to 42 wherein the viral disorder is selected from human cytomegalovirus (HCMV), herpes simplex virus type 1 (HSV-1), human immunodeficiency virus type 1 (HIV-1), and varicella zoster virus (VZV).
44. Use of a compound according to any one of claims 1 to 34 for inhibiting a protein kinase.
45. Use according to claim 44 wherein said protein kinase is a cyclin dependent kinase.
46. Use according to claim 45 wherein said protein dependent kinase is selected from CDK2, CDK7, CDK8 and CDK9.
47. A method of treating a proliferative disease, said method comprising administering to a mammal a therapeutically effective amount of a compound according to any one of claims 1 to 34.
48. A method of treating a viral disorder, said method comprising administering to a mammal a therapeutically effective amount of a compound of formula Ia, or a pharmaceutically acceptable salt thereof,



wherein

- (A) one of X^1 and X^2 is S, and the other of X^1 and X^2 is N;
 "a" is a single bond; and
 "b", "c", "d", "e" and "f" are single or double bonds so as to form a thiazolyl ring;
 R^2 is independently as defined below for R^1 , R^{3-8} ; or
- (B) one of X^1 and X^2 is S, and the other of X^1 and X^2 is NR^{17} ;
 "a" and "d" are each double bonds; and
 "b", "c", "e" and "f" are each single bonds;
 R^2 is oxo;
 R^{17} is H or alkyl;

where:

Z is NH, NHCO, NHSO₂, NHCH₂, CH₂, CH₂CH₂, CH=CH, SO₂, or SO;

R^1 , R^2 , R^3 , R^4 , R^5 , R^6 , R^7 and R^8 are each independently H, alkyl, alkyl- R^9 , aryl, aryl- R^9 , aralkyl, aralkyl- R^9 , halogeno, NO₂, CN, OH, O-alkyl, COR⁹, COOR⁹, O-aryl, O- R^9 , NH₂, NH-alkyl, NH-aryl, N-(alkyl)₂, N-(aryl)₂, N-(alkyl)(aryl), NH- R^9 , N-(R^9)(R^{10}), N-(alkyl)(R^9), N-(aryl)(R^9), COOH, CONH₂, CONH-alkyl, CONH-aryl, CON-(alkyl)(R^9), CON(aryl)(R^9), CONH- R^9 , CON-(R^9)(R^{10}), SO₃H, SO₂-alkyl, SO₂-alkyl- R^9 , SO₂-aryl, SO₂-aryl- R^9 , SO₂NH₂, SO₂NH- R^9 , SO₂N-(R^9)(R^{10}), CF₃, CO-alkyl, CO-alkyl- R^9 , CO-aryl, CO-aryl- R^9 or R^{11} , wherein alkyl, aryl, aralkyl groups may be further substituted with one or more groups selected from halogeno, NO₂, OH, O-methyl, NH₂, COOH, CONH₂ and CF₃;

wherein at least one of R^1 , R^2 , R^3 , R^4 , R^5 , R^6 , R^7 and R^8 is an R^9 or R^{10} -containing group, or is R^{11} ;

R^9 and R^{10} are each independently solubilising groups selected from:

- (i) - a mono-, di- or polyhydroxylated alicyclic group;
- a di- or polyhydroxylated aliphatic or aromatic group;
- a carbohydrate derivative;

121

- an O- and/or S-containing heterocyclic group optionally substituted by one or more hydroxyl groups;
 - an aliphatic or aromatic group containing a carboxamide, sulfoxide, sulfone, or sulfonamide function; or
 - a halogenated alkylcarbonyl group;
- (ii) COOH , SO_3H , OSO_3H , PO_3H_2 , or OPO_3H_2 ;
- (iii) Y, where Y is selected from an alicyclic, aromatic, or heterocyclic group comprising one or more of the functions $=\text{N}-$, $-\text{O}-$, $-\text{NH}_2$, $-\text{NH}-$, a quarternary amine salt, guanidine, and amidine, where Y is optionally substituted by one or more substituents selected from:
- SO_2 -alkyl;
 - alkyl optionally substituted by one or more OH groups;
 - CO -alkyl;
 - aralkyl;
 - COO -alkyl; and
 - an ether group optionally substituted by one or more OH groups; and
- where Y is other than pyridinyl;
- (iv) a natural or unnatural amino acid, a peptide or a peptide derivative;

each R^{11} is a solubilising group as defined for R^9 and R^{10} in (i) or (iv) above; or is selected from:

- (v) OSO_3H , PO_3H_2 , or OPO_3H_2 ;
- (vi) Y as defined above, but excluding guanidine and quarternary amine salts;
- (vii) $\text{NHCO}(\text{CH}_2)_m[\text{NHCO}(\text{CH}_2)_{m'}]_p[\text{NHCO}(\text{CH}_2)_{m''}]_q\text{Y}$ or $\text{NHCO}(\text{CH}_2)_t\text{NH}(\text{CH}_2)_{t'}\text{Y}$ where p and q are each 0 or 1, and m, m', m'', t and t' are each an integer from 1 to 10; and
- (viii) $(\text{CH}_2)_n\text{NR}^{14}\text{COR}^{12}$, $(\text{CH}_2)_n\text{NR}^{15}\text{SO}_2\text{R}^{13}$, or SO_2R^{16} , where R^{12} , R^{13} and R^{16} are each alkyl groups optionally comprising one or more heteroatoms, and which are substituted by one or more substituents selected from OH, NH_2 , halogen and NO_2 ,

122

R^{14} and R^{15} are each independently H or alkyl, and n and n' are each independently 0, 1, 2, or 3;

- (ix) an ether or polyether optionally substituted by one or more hydroxyl groups or one or more Y groups;
- (x) $(CH_2)_rNH_2$; where r is 0, 1, 2, or 3;
- (xi) $(CH_2)_{r'}OH$; where r' is 0, 1, 2, or 3.

49. Use of a compound of formula I as defined in any one of claims 1 to 34 in an assay for identifying further candidate compounds capable of inhibiting one or more of a cyclin dependent kinase, GSK and a PLK enzyme.

50. Use according to claim 49 wherein said assay is a competitive binding assay.

51. Use according to claim 50 wherein said competitive binding assay comprises contacting a compound of formula I as defined in any one of claims 1 to 34 with a cyclin dependent kinase and a candidate compound and detecting any change in the interaction between the compound of formula I and the cyclin dependent kinase.

52. Use according to claim 36 wherein the proliferative disorder is glomerulonephritis, rheumatoid arthritis, psoriasis or chronic obstructive pulmonary disorder.

53. Use of a compound of formula I or Ia, or a pharmaceutically acceptable salt thereof, in the preparation of a medicament for treating a CNS disorder.

54. Use according to claim 53 wherein the CNS disorder is Alzheimer's disease or bipolar disorder.

55. Use of a compound of formula I or Ia, or a pharmaceutically acceptable salt thereof, in the preparation of a medicament for treating alopecia.

56. Use of a compound of formula I or Ia, or a pharmaceutically acceptable salt thereof, in the preparation of a medicament for treating a stroke.
57. Use according to any one of claims 36, 37 or 52 to 56 wherein the compound is administered in an amount sufficient to inhibit at least one PLK enzyme.
58. Use according to claim 57 wherein the PLK enzyme is PLK1.
59. Use according to any one of claims 36, 37 or 52 to 56 wherein the compound is administered in an amount sufficient to inhibit at least one CDK enzyme.
60. Use according to claim 59 wherein the CDK enzyme is CDK1, CDK2, CDK3, CDK4, CDK6, CDK7, CDK8 and/or CDK9.
61. Use according to any one of claims 36, 37 or 52 to 56 wherein the compound is administered in an amount sufficient to inhibit aurora kinase.
62. Use of a compound of formula I or Ia, or a pharmaceutically acceptable salt thereof, in the preparation of a medicament for treating diabetes.
63. Use according to claim 62 wherein the diabetes is Type II diabetes.
64. Use according to any one of claims 62 or 63 wherein the compound is administered in an amount sufficient to inhibit GSK.
65. Use according to claim 64 wherein the compound is administered in an amount sufficient to inhibit GSK3 β .

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/GB 03/04973

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D417/04 A61K31/506 A61P35/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

CHEM ABS Data, PAJ

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 97 19065 A (CELLTECH) 29 May 1997 (1997-05-29) claims; examples 8,9,26,30	1-5, 35-39
X	WO 01 72745 A (CYCLACEL) 4 October 2001 (2001-10-04) claims 1-4,17-20; examples	1-5, 35-39
P,X	WO 03 29248 A (CYCLACEL) 10 April 2003 (2003-04-10) page 27; claims; examples 4,18	1-5,14, 26,35-39

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

Z document member of the same patent family

Date of the actual completion of the international search

12 February 2004

Date of mailing of the international search report

27/02/2004

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Francois, J

INTERNATIONAL SEARCH REPORT

International application No.
PCT/GB 03/04973

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

Although claims 47,48 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/GB 03/04973

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9719065 A	29-05-1997	AU 7631496 A	11-06-1997
		DE 69627179 D1	08-05-2003
		DE 69627179 T2	29-01-2004
		EP 0862560 A1	09-09-1998
		ES 2195020 T3	01-12-2003
		WO 9719065 A1	29-05-1997
		US 6235746 B1	22-05-2001
		US 5958935 A	28-09-1999
WO 0172745 A	04-10-2001	AU 4262901 A	08-10-2001
		CA 2401748 A1	04-10-2001
		CN 1420884 T	28-05-2003
		EP 1274705 A1	15-01-2003
		WO 0172745 A1	04-10-2001
		GB 2361236 A ,B	17-10-2001
		HU 0300382 A2	28-06-2003
		JP 2003528872 T	30-09-2003
		US 2003149057 A1	07-08-2003
		US 2002019404 A1	14-02-2002
WO 0329248 A	10-04-2003	WO 03029248 A1	10-04-2003